arXiv:0901.4471v4 [math-ph] 19 Jul 2010

Classification of two and three dimensional Lie super-bialgebras

A. Eghbali [*], A. Rezaei-Aghdam [†] and F. Heidarpour
Department of Physics, Faculty of science, Azarbaijan University
of Tarbiat Moallem , 53714-161, Tabriz, Iran

July 20, 2010

Abstract

Using adjoint representation of Lie superalgebras, we write the matrix form of super Jacobi and mixed super Jacobi identities of Lie super-bialgebras. Then through direct calculations of these identities and use of automorphism supergroups of two and three dimensional Lie superalgebras, we obtain and classify all two and three dimensional Lie super-bialgebras.

---

[*] e-mail: a.eghbali@azaruniv.edu
[†] Corresponding author. e-mail: rezaei-a@azaruniv.edu

1 Introduction

Lie bialgebras were first introduced from the mathematical point of view, by Drinfel'd as algebraic structures and classical limit of underlying quantized enveloping algebras (*quantum groups*) [1]. In particular, every deformation of a universal enveloping algebra induces a Lie bialgebra structure on the underlying Lie algebra. Conversely it has been shown in [2] that each Lie bialgebra admits quantization. So the classification of Lie bialgebras can be seen as the first step in the classification of quantum groups. Many interesting examples of Lie bialgebras based on complex semisimple Lie algebras have been given in Derinfel'd [1]. A complete classification of Lie bialgebras with reduction double was given in [3]. However a classification of Lie bialgebras is out of reach, just for similar reasons as for the Lie algebra classification. In the non-semisimple case only a bunch of low dimensional examples has been thoroughly studied [4, 5, 6, 7]. On the other hand, from the physical point of view, the theory of classical integrable system naturally relates to the geometry and representation theory of Poisson-Lie groups and the corresponding Lie bialgebras and their classical r-matrices (see for example [8]). In the same way, Lie super-bialgebras [9], as the underlying symmetry algebras, play an important role in the integrable structure of AdS/CFT correspondence [10]. In this way and by considering that there is a universal quantization for Lie super-bialgebras [11], one can assign an important role to the classification of Lie super-bialgebras (especially low dimensional Lie super-bialgebras) from both physical and mathematical point of view. There are distinguished and nonsystematic ways for obtaining low dimensional Lie super-bialgebras (see for example [12, 13, 14]). In this paper, using adjoint representation of Lie superalgebras, we give a systematic way for obtaining and classification of low dimensional Lie super-bialgebras. We apply this method to the classification of two and three dimensional Lie super-bialgebras.

The paper is organized as follows. In section two, we give basic definitions and notations that are used throughout the paper. The systematic way of classification of Lie super-bialgebras by using matrix form of super Jacobi (sJ) and mixed super Jacobi (msJ) identities of Lie super-bialgebras is described in section three. A list of two and three dimensional Lie superalgebras of [15] is offered in section 4. The obtained automorphism Lie supergroups of these Lie superalgebras are also presented in section 4 . Then, using the method mentioned in section two, we classify all 48 two and three dimensional Lie super-bialgebras in section five. The details of calculations are explained using an example. The concluding section discusses some remarks. Certain properties of tensors and supermatrices are brought in the appendix A. Solutions of ($sJ - msJ$) identities and their isomorphism matrices are give in the appendix B.

2 Basic definitions and notations

In this paper we use DeWitt notation for supervector spaces, supermatrices,...[16]. Some of definitions and related notations are brought in appendix A.

Definition: A *Lie superalgebra* $\mathbf{g}$ is a graded vector space $\mathbf{g} = \mathbf{g}_B \bigoplus \mathbf{g}_F$ with gradings; $grade(\mathbf{g}_B) = 0$, $grade(\mathbf{g}_F) = 1$; such that Lie bracket satisfies the super antisymmetric and super Jacobi identities, i.e. in a graded basis $\{X_i\}$ of $\mathbf{g}$ if we put[1]

$$[X_i, X_j] = f^k{}_{ij} X_k, \tag{1}$$

then

$$(-1)^{i(j+k)} f^m{}_{jl} f^l{}_{ki} + f^m{}_{il} f^l{}_{jk} + (-1)^{k(i+j)} f^m{}_{kl} f^l{}_{ij} = 0, \tag{2}$$

so that

$$f^k{}_{ij} = -(-1)^{ij} f^k{}_{ji}. \tag{3}$$

Note that, in the standard basis, $f^B{}_{BB}$ and $f^F{}_{BF}$ are real c-numbers and $f^B{}_{FF}$ are pure imaginary c-numbers and other components of structure constants $f^i{}_{jk}$ are zero [16], i.e. we have

$$f^k{}_{ij} = 0, \qquad if \quad grade(i) + grade(j) \neq grade(k) \quad (mod 2). \tag{4}$$

---

[1] Note that the bracket of one boson with one boson or one fermion is usual commutator but for one fermion with one fermion is anticommutator. Furthermore we identify grading of indices by the same indices in the power of (-1), for example $grading(i) \equiv i$; this is the notation that DeWitt applied in his book [16] .

Let $\mathbf{g}$ be a finite-dimensional Lie superalgebra and $\mathbf{g}^*$ be its dual vector space and let $(\ ,\ )$ be the canonical pairing on $\mathbf{g}^* \bigoplus \mathbf{g}$.

Definition: A *Lie super-bialgebra* structure on a Lie superalgebra $\mathbf{g}$ is a super skew-symmetric linear map $\delta : \mathbf{g} \longrightarrow \mathbf{g} \otimes \mathbf{g}$ (*the super cocommutator*) so that[2] [9]

1)δ is a super one-cocycle, i.e.

$$\delta([X,Y]) = (ad_X \otimes I + I \otimes ad_X)\delta(Y) - (-1)^{|X||Y|}(ad_Y \otimes I + I \otimes ad_Y)\delta(X) \qquad \forall X, Y \in \mathbf{g}, \tag{5}$$

2) the dual map ${}^t\delta : \mathbf{g}^* \otimes \mathbf{g}^* \to \mathbf{g}^*$ is a Lie superbracket on $\mathbf{g}^*$, i.e.

$$(\xi \otimes \eta, \delta(X)) = ({}^t\delta(\xi \otimes \eta), X) = ([\xi, \eta]_*, X) \qquad \forall X \in \mathbf{g};\ \ \xi, \eta \in \mathbf{g}^*. \tag{6}$$

The Lie super-bialgebra defined in this way will be denoted by $(\mathbf{g}, \mathbf{g}^*)$ or $(\mathbf{g}, \delta)$.

Proposition: Super one-cocycles δ and δ' of the Lie superalgebra $\mathbf{g}$ are said to be *equivalent* if there exists an automorphism A of $\mathbf{g}$ such that

$$\delta' = (A \otimes A) \circ \delta \circ A^{-1}. \tag{7}$$

In this case the two Lie super-bialgebras namely $(\mathbf{g}, \delta)$ and $(\mathbf{g}, \delta')$, are equivalent [7].

Definition: A *Manin* super triple [9] is a triple of Lie superalgebras $(\mathcal{D}, \mathbf{g}, \tilde{\mathbf{g}})$ together with a non-degenerate ad-invariant super symmetric bilinear form $<\ ,\ >$ on $\mathcal{D}$ such that

1) $\mathbf{g}$ and $\tilde{\mathbf{g}}$ are Lie sub-superalgebras of $\mathcal{D}$,

2) $\mathcal{D} = \mathbf{g} \bigoplus \tilde{\mathbf{g}}$ as a supervector space,

3) $\mathbf{g}$ and $\tilde{\mathbf{g}}$ are isotropic with respect to $<,>$, i.e.

$$< X_i, X_j > = < \tilde{X}^i, \tilde{X}^j > = 0, \qquad \delta_i{}^j = < X_i, \tilde{X}^j > = (-1)^{ij} < \tilde{X}^j, X_i > = (-1)^{ij} \delta^j{}_i, \tag{8}$$

where $\{X_i\}$ and $\{\tilde{X}^i\}$ are basis of Lie superalgebras $\mathbf{g}$ and $\tilde{\mathbf{g}}$, respectively. Note that in the above relation $\delta^j{}_i$ is the ordinary delta function. There is a one-to-one correspondence between Lie super-bialgebra $(\mathbf{g}, \mathbf{g}^*)$ and Manin super triple $(\mathcal{D}, \mathbf{g}, \tilde{\mathbf{g}})$ with $\mathbf{g}^* = \tilde{\mathbf{g}}$ [9]. If we choose the structure constants of Lie superalgebras $\mathbf{g}$ and $\tilde{\mathbf{g}}$ as

$$[X_i, X_j] = f^k{}_{ij} X_k, \qquad [\tilde{X}^i, \tilde{X}^j] = \tilde{f}^{ij}{}_k \tilde{X}^k, \tag{9}$$

then ad-invariance of the bilinear form $<,>$ on $\mathcal{D} = \mathbf{g} \bigoplus \tilde{\mathbf{g}}$ implies that

$$[X_i, \tilde{X}^j] = (-1)^j \tilde{f}^{jk}{}_i X_k + (-1)^i f^j{}_{ki} \tilde{X}^k. \tag{10}$$

Clearly, using the equations (6) and (9) we have[3]

$$\delta(X_i) = (-1)^{jk} \tilde{f}^{jk}{}_i X_j \otimes X_k. \tag{11}$$

As a result of applying this relation to the super one-cocycle condition (5), the super Jacobi identities (2) for the dual Lie superalgabra and the following mixed super Jacobi identities are obtained[4]

$$f^m{}_{jk} \tilde{f}^{il}{}_m = f^i{}_{mk} \tilde{f}^{ml}{}_j + f^l{}_{jm} \tilde{f}^{im}{}_k + (-1)^{jl} f^i{}_{jm} \tilde{f}^{ml}{}_k + (-1)^{ik} f^l{}_{mk} \tilde{f}^{im}{}_j. \tag{12}$$

---

[2] Here $|X|(|Y|)$ indicates the grading of $X(Y)$.

[3] Note that the appearance of $(-1)^{jk}$ in this relation is due to the definition of natural inner product between $\mathbf{g} \otimes \mathbf{g}$ and $\mathbf{g}^* \otimes \mathbf{g}^*$ as $(\tilde{X}^i \otimes \tilde{X}^j, X_k \otimes X_l) = (-1)^{jk} \delta^i{}_k \delta^j{}_l$.

[4] This relation can also be obtained from super Jacobi identity of $\mathcal{D}$.

3 Calculation of Lie super-bialgebras using adjoint representation

As discussed in the introduction, by use of super Jacobi and mixed super Jacobi identities we classify two and three dimensional Lie super-bialgebras. This method was first used to obtain three dimensional Lie bialgebras in [4]. Because of tensorial form of super Jacobi and mixed super Jacobi identities, working with them is not so easy and we suggest writing these equations as matrix forms using the following adjoint representations for Lie superalgebras $\mathbf{g}$ and $\tilde{\mathbf{g}}$

$$(\tilde{\mathcal{X}}^i)^j{}_k = -\tilde{f}^{ij}{}_k, \qquad (\mathcal{Y}^i)\,{}_{jk} = -f^i{}_{jk}. \tag{13}$$

Then the matrix forms of super Jacobi identities (2) for dual Lie superalgebra $\tilde{\mathbf{g}}$ and mixed super Jacobi identities (12) become as follows respectively:

$$(\tilde{\mathcal{X}}^i)^j{}_k \tilde{\mathcal{X}}^k - \tilde{\mathcal{X}}^j \tilde{\mathcal{X}}^i + (-1)^{ij} \tilde{\mathcal{X}}^i \tilde{\mathcal{X}}^j = 0, \tag{14}$$

$$(\tilde{\mathcal{X}}^i)^j{}_l\, \mathcal{Y}^l = -(-1)^k \tilde{\mathcal{X}}^{st\ j}\, \mathcal{Y}^i + \mathcal{Y}^j \tilde{\mathcal{X}}^i - (-1)^{ij} \mathcal{Y}^i \tilde{\mathcal{X}}^j + (-1)^{k+ij} \tilde{\mathcal{X}}^{st\ i}\, \mathcal{Y}^j. \tag{15}$$

Note that in the above relations we use the right indices for the matrix elements and index k represents the column of matrix $\tilde{\mathcal{X}}^{st}$.

Having the structure constants of the Lie superalgebra $\mathbf{g}$, we solve the matrix equations (14) and (15) in order to obtain the structure constants of the dual Lie superalgebras $\tilde{\mathbf{g}}$ so that $(\mathbf{g}, \tilde{\mathbf{g}})$ is Lie super-bialgebra. We assume that, we have classification and tabulation of the Lie superalgebras $\mathbf{g}$ (for example two and three dimensional Lie superalgebras for this paper)[5]. We fulfill this work in the following three steps using Maple.

Step 1) With the solution of matrix equations (14) and (15) for obtaining matrices $\tilde{\mathcal{X}}^i$, some structure constants of $\tilde{\mathbf{g}}$ are obtained zero, and some unknown or in terms of each other. In order to know whether $\tilde{\mathbf{g}}$ is one of the Lie superalgebras of table or isomorphic to those, we must use the following isomorphic relation between obtained Lie superalgebras $\tilde{\mathbf{g}}$ and one of the Lie superalgebras of table, e.g. $\mathbf{g}'$. Applying the transformation (37) for a change of basis $\tilde{\mathbf{g}}$ we have

$$\tilde{X}'^{\ i} = C^i{}_j \tilde{X}^j, \qquad [\tilde{X}'^{\ i}, \tilde{X}'^{\ j}] = \tilde{f}'^{\ ij}_{\ k} \tilde{X}'^{\ k}, \tag{16}$$

then the following matrix equations for isomorphism

$$(-1)^{i(j+l)}\, C\, (C^i{}_k\, \tilde{\mathcal{X}}^k_{(\tilde{\mathbf{g}})}) = \mathcal{X}^i_{(\mathbf{g}')}\, C, \tag{17}$$

are obtained, where the indices j and l corresponds to the row and column of C in the left hand side of (17). In the above matrix equations $\mathcal{X}^i_{(\mathbf{g}')}$ are adjoint matrices of known Lie superalgebra $\mathbf{g}'$ of the classification table. Solving (17) with the condition $sdetC \neq 0$ we obtain some extra conditions on $\tilde{f}^{kl}_{(\tilde{\mathbf{g}})\ m}$'s that were obtained from (14) and (15).

Step 2) As the second step we transform Lie super-bialgebra $(\mathbf{g}, \tilde{\mathbf{g}})$ (where in the Lie superalgebra $\tilde{\mathbf{g}}$ we impose extra conditions obtained in the step one) to Lie super bialgebra $(\mathbf{g}, \mathbf{g}'.i)$ (where $\mathbf{g}'.i$ is isomorphic as Lie superalgebra to $\mathbf{g}'$) with an automorphism of the Lie superalgebra $\mathbf{g}$. As the inner product (8) is invariant we have $A^{-st} : \tilde{\mathbf{g}} \longrightarrow \mathbf{g}'.i$,

$$X'_i = (-1)^k A_i{}^k X_k, \qquad \tilde{X}'^{\ j} = (A^{-st})^j{}_l \tilde{X}^l, \qquad < X'_i, \tilde{X}'^{\ j} >= \delta_i{}^j, \tag{18}$$

where A^{-st} is superinverse supertranspose of every matrix $A \in Aut(\mathbf{g})$ [6]. Thus we have the following transformation relation for the map A^{-st}

$$(-1)^{k(j+l)} (A^{-st})^i{}_k \tilde{f}^{kl}_{(\tilde{\mathbf{g}})\ m} (A^{-st})^j{}_l = f^{ij}{}_{(\mathbf{g}'.\mathbf{i})\ n} (A^{-st})^n{}_m. \tag{19}$$

Now, for obtaining Lie super-bialgebras $(\mathbf{g}, \mathbf{g}'.i)$, we must find Lie superalgebras $\mathbf{g}'.i$ or transformations $B : \mathbf{g}' \longrightarrow \mathbf{g}'.i$ such that

$$(-1)^{k(j+l)} B^i{}_k f^{kl}{}_{(\mathbf{g}')\ m} B^j{}_l = f^{ij}{}_{(\mathbf{g}'.\mathbf{i})\ n} B^n{}_m. \tag{20}$$

---

[5] Here we explain the method in general and one can apply it for classification of other low dimensional Lie super-bialgebras [17].

[6] $Aut(\mathbf{g})$ is the automorphism supergroup of the Lie superalgebra $\mathbf{g}$

For this purpose, it is enough to omit ${f^{ij}}_{(\mathbf{g'.i})\ n}$ between (19) and (20). Then we will have the following matrix equation for B [7]

$$(A^{-st})^i{}_m \tilde{\mathcal{X}}^{st\ m}_{(\tilde{\mathbf{g}})} A^{-1} = (-1)^{i(l+j)} (B^{st} A)^{-1} (B^i{}_k {\mathcal{X}^{st\ k}}_{(\mathbf{g'})}) B^{st}, \tag{21}$$

where indices l and j are row and column of B^{st} in the right hand side of the above relation. Now by solving (21) we obtain the general form of matrix B with the condition $sdetB \neq 0$. In solving (21) one can obtain conditions on elements of matrix A, yet we must only consider those conditions under which we have $sdetA \neq 0$ and matrices A, B and A^{-st} have the general transformation matrix form (38).

Step 3) Having solved (21), we obtain the general form of the matrix B so that its elements are written in terms of the elements of matrices A, C and structure constants $\tilde{f}^{ij}_{(\tilde{\mathbf{g}})\ k}$. Now with substituting B in (20), we obtain structure constants ${f^{ij}}_{(\mathbf{g'.i})\ n}$ of the Lie superalgebra $\mathbf{g'}.i$ in terms of elements of matrices A and C and some $\tilde{f}^{ij}_{(\tilde{\mathbf{g}})\ k}$. Then we check whether it is possible to equalize the structure constants ${f^{ij}}_{(\mathbf{g'.i})\ n}$ with each other and with ± 1 or not so as to remark $sdetB \neq 0$, $sdetA \neq 0$ and $sdetC \neq 0$. In this way, we obtain isomorphism matrices B_1, B_2,... . As B_i^{st} must be in the form of transformation matrices (38), we obtain conditions on the matrices B_i. The reason is that if $(\mathbf{g}, \mathbf{g'}.i)$ is Lie super-bialgebra, then $(\mathbf{g'}, \mathbf{g}.i)$ will be Lie super-bialgebras with $B_i^{st} : \mathbf{g} \longrightarrow \mathbf{g}.i$. Note that in obtaining B_is we impose the condition $B{B_i}^{-1} \in Aut^{st}(\mathbf{g})$ [8]; if this condition is not satisfied then we can not impose it on the structure constants because B and B_i are not equivalent (see bellow).

Now using isomorphism matrices B_1, B_2,... , we can obtain Lie super-bialgebras $(\mathbf{g}, \mathbf{g'}.i)$, $(\mathbf{g}, \mathbf{g'}.ii)$,... . On the other hand, there is the question: which of these Lie super-bialgebras are equivalent? In order to answer this question, we use the matrix form of the relation (7). Consider the two Lie super-bialgebras $(\mathbf{g}, \mathbf{g'}.i)$, $(\mathbf{g}, \mathbf{g'}.ii)$; then using [9]

$$A(X_i) = (-1)^j {A_i}^j X_j, \tag{22}$$

the relation (7) will have the following matrix form

$$(-1)^{i(j+l)} A^{st} ((A^{st})^i{}_k {\mathcal{X}_{(\mathbf{g'}.i)}}^k) = {\mathcal{X}_{(\mathbf{g'}.ii)}}^i A^{st}. \tag{23}$$

On the other hand, the transformation matrix between $\mathbf{g'}.i$ and $\mathbf{g'}.ii$ is $B_2 B_1^{-1}$ if $B_1 : \mathbf{g'} \longrightarrow \mathbf{g'}.i$ and $B_2 : \mathbf{g'} \longrightarrow \mathbf{g'}.ii$; then we have

$$(-1)^{i(j+l)} (B_2 B_1^{-1}) ((B_2 B_1^{-1})^i{}_k {\mathcal{X}_{(\mathbf{g'}.i)}}^k) = {\mathcal{X}_{(\mathbf{g'}.ii)}}^i (B_2 B_1^{-1}). \tag{24}$$

A comparison of (24) with (23) reveals that if $B_2 B_1^{-1} \in A^{st}$ holds, then the Lie super-bialgebras $(\mathbf{g}, \mathbf{g'}.i)$ and $(\mathbf{g}, \mathbf{g'}.ii)$ are equivalent. In this way, we obtain nonequivalent class of B_is and we consider only one element of this class. Thus, we obtain and classify all Lie super-bialgebras. In the next section, we apply this formulation to two and three dimensional Lie superalgebras.

4 Two and three dimensional Lie superalgebras and their automorphism supergroups

In this section, we use the classification of two and three dimensional Lie superalgebras listed in [15]. In this classification, Lie superalgebras are divided into two types: trivial and nontrivial Lie superalgebras for which the commutations of fermion-fermion is zero or nonzero respectively (as we use DeWitt notation here, the structure constant C^B_{FF} must be pure imaginary). The results have been presented in tables 1 and 2. As the tables $(m, n-m)$ indicate the Lie superalgebras have $m, \{X_1, ..., X_m\}$ bosonic and $n-m, \{X_{m+1}, ..., X_n\}$ fermionic generators.

---

[7] Note that for A^{-st} and A^{-1} in (21) we use row and column indices on the right hand sides.

[8] $Aut^t(\mathbf{g})$ is the supertranspose of $Aut(\mathbf{g})$.

[9] Note that the effect of A is linear and even i.e.

$$A({a_i}^j X_j) = {a_i}^j A(X_j).$$

Table 1 : Trivial Lie superalgebras

Type	**g**	Bosonic bases	Fermionic bases	Commutation relations	Comments
$(1,1)$	B	X_1	X_2	$[X_1,X_2]=X_2$	
$(2,1)$	C_p^1	X_1,X_2	X_3	$[X_1,X_2]=X_2,\ [X_1,X_3]=pX_3$	$p\neq 0$
$(1,2)$	C_p^2	X_1	X_2,X_3	$[X_1,X_2]=X_2,\ [X_1,X_3]=pX_3$	$0<\lvert p\rvert\leq 1$
	C^3	X_1	X_2,X_3	$[X_1,X_3]=X_2$	Nilpotent
	C^4	X_1	X_2,X_3	$[X_1,X_2]=X_2,\ [X_1,X_3]=X_2+X_3$	
	C_p^5	X_1	X_2,X_3	$[X_1,X_2]=pX_2-X_3,\ [X_1,X_3]=X_2+pX_3$	$p\geq 0$

Table 2 : Nontrivial Lie superalgebras[10].

Type	**g**	Bosonic bases	Fermionic bases	Commutation relations	Comments
$(1,1)$	$(A_{1,1}+A)$	X_1	X_2	$\{X_2,X_2\}=iX_1$	
$(2,1)$	$C_{\frac{1}{2}}^1$	X_1,X_2	X_3	$[X_1,X_2]=X_2,\ [X_1,X_3]=\frac{1}{2}X_3,\ \{X_3,X_3\}=iX_2$	
$(1,2)$	$(A_{1,1}+2A)^1$	X_1	X_2,X_3	$\{X_2,X_2\}=iX_1,\ \{X_3,X_3\}=iX_1$	Nilpotent
	$(A_{1,1}+2A)^2$	X_1	X_2,X_3	$\{X_2,X_2\}=iX_1,\ \{X_3,X_3\}=-iX_1$	Nilpotent

One can check whether super Jacobi identities for the above Lie superalgebras are satisfied. As mentioned in section 3 for obtaining dual Lie superalgebras we need automorphism supergroups of Lie superalgebras. In order to calculate the automorphism supergroups of two and three dimensional Lie superalgebras, we use the following transformation:

$$X'_i = (-1)^j\ A_i{}^j X_j, \qquad\qquad [X'_i, X'_j] = f^k{}_{ij} X'_k, \tag{25}$$

thus we have the following matrix equation for elements of automorphism supergroups

$$(-1)^{ij+mk} A\mathcal{Y}^k A^{st} = \mathcal{Y}^e A_e{}^k, \tag{26}$$

where index j is the column of matrix A and indices i and m are row and column of A^{st} in the left hand side. Using (26) with the condition $sdetA \neq 0$ and imposing the condition that A must be in the form (38), we obtain table 3 for automorphism supergroups

---

[10] Note that the superalgebra A is a one dimensional abelian Lie superalgebras with one fermionic generator and superalgebra $A_{1,1}$ is its bosonization.

Table 3 : Automorphism supergroups of the two and three dimensional Lie superalgebras

$\mathbf{g}$	Automorphism supergroups	comments
B	$\begin{pmatrix} 1 & 0 \\ 0 & b \end{pmatrix}$	$b \in \Re - \{0\}$
$(A_{1,1}+A)$	$\begin{pmatrix} a^2 & 0 \\ 0 & a \end{pmatrix}$	$a \in \Re - \{0\}$
C_p^1	$\begin{pmatrix} 1 & a & 0 \\ 0 & c & 0 \\ 0 & 0 & d \end{pmatrix}$	$p \in \Re - \{0\}, \quad c, d \in \Re - \{0\}, \ a \in \Re$
$C_{\frac{1}{2}}^1$	$\begin{pmatrix} 1 & a & 0 \\ 0 & b^2 & 0 \\ 0 & 0 & b \end{pmatrix}$	$b \in \Re - \{0\}$, $a \in \Re$
C_1^2	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & b \\ 0 & a & d \end{pmatrix}$	$ab - cd \neq 0$
C_p^2	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & d \end{pmatrix}$	$c, d \in \Re - \{0\}, \ \ p \in [-1, 1] - \{0\}$
C^3	$\begin{pmatrix} a & 0 & 0 \\ 0 & ad & 0 \\ 0 & e & d \end{pmatrix}$	$a, d \in \Re - \{0\}$, $e \in \Re$
C^4	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & 0 \\ 0 & d & c \end{pmatrix}$	$c \in \Re - \{0\}, \ \ d \in \Re$
C_p^5	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & -d \\ 0 & d & c \end{pmatrix}$	$p \geq 0, \ \ c \in \Re - \{0\}$ or $d \in \Re - \{0\}$
$(A_{1,1}+2A)^1$	$\begin{pmatrix} b^2+c^2 & 0 & 0 \\ 0 & b & -c \\ 0 & c & b \end{pmatrix}$	$b \in \Re - \{0\}$ or $c \in \Re - \{0\}$
$(A_{1,1}+2A)^2$	$\begin{pmatrix} b^2-c^2 & 0 & 0 \\ 0 & b & c \\ 0 & c & b \end{pmatrix}$	$b \in \Re - \{0\}$ or $c \in \Re - \{0\}$

5 Two and three dimensional Lie super-bialgebras

Using the methods discussed in section 3, we can classify two and three dimensional Lie super-bialgebras. We have applied Maple 9 and obtained 48 Lie super-bialgebras. These have been listed in the following tables 4-7:

Table 4 : Two dimensional Lie super-bialgebras of the type $(1,1)$

$\mathbf{g}$	$\tilde{\mathbf{g}}$	(Anti) Commutation relations of $\tilde{\mathbf{g}}$
$(A_{1,1}+A)$	$I_{(1,1)}$	
B	$I_{(1,1)}$	
	$(A_{1,1}+A)$	$\{\tilde{X}^2, \tilde{X}^2\} = i\tilde{X}^1$
	$(A_{1,1}+A).i$	$\{\tilde{X}^2, \tilde{X}^2\} = -i\tilde{X}^1$

Table 5: Three dimensional Lie super-bialgebras of the type (2 ,1).

$\mathbf{g}$	$\tilde{\mathbf{g}}$	(Anti) Commutation relations of $\tilde{\mathbf{g}}$	Comments
C_p^1	$I_{(2,1)}$		
	$C^1_{-p}.i$	$[\tilde{X}^1,\tilde{X}^2]=\tilde{X}^1,\quad [\tilde{X}^2,\tilde{X}^3]=p\tilde{X}^3$	$p\in\Re-\{0\}$
$C^1_{\frac{1}{2}}$	$I_{(2,1)}$		
	$C^1_p.i\|_{p=-\frac{1}{2}}$	$[\tilde{X}^1,\tilde{X}^2]=\tilde{X}^1,\quad [\tilde{X}^2,\tilde{X}^3]=\frac{1}{2}\tilde{X}^3$	
	$C^1_p.ii\|_{p=-\frac{1}{2}}$	$[\tilde{X}^1,\tilde{X}^2]=-\tilde{X}^1,\quad [\tilde{X}^2,\tilde{X}^3]=-\frac{1}{2}\tilde{X}^3$	
	$C^1_{\frac{1}{2}}.i$	$[\tilde{X}^1,\tilde{X}^2]=-\tilde{X}^1,\quad [\tilde{X}^2,\tilde{X}^3]=\frac{1}{2}\tilde{X}^3,\quad \{\tilde{X}^3,\tilde{X}^3\}=i\tilde{X}^1$	
	$C^1_{\frac{1}{2}}.ii$	$[\tilde{X}^1,\tilde{X}^2]=\tilde{X}^1,\quad [\tilde{X}^2,\tilde{X}^3]=-\frac{1}{2}\tilde{X}^3,\quad \{\tilde{X}^3,\tilde{X}^3\}=-i\tilde{X}^1$	
	$C^1_{\frac{1}{2},k}$	$[\tilde{X}^1,\tilde{X}^2]=-k\tilde{X}^2,\ [\tilde{X}^1,\tilde{X}^3]=-\frac{k}{2}\tilde{X}^3,\ \{\tilde{X}^3,\tilde{X}^3\}=k\tilde{X}^2$	$k\in\Re-\{0\}$

Table 6: Three dimensional Lie super-bialgebras of the type (1 ,2)

$\mathbf{g}$	$\tilde{\mathbf{g}}$	Comments
C_1^2	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{\epsilon,0,\epsilon}$	
	$(A_{1,1}+2A)^2_{\epsilon,0,-\epsilon}$	
C_p^2	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{\epsilon,k,\epsilon}$	$-1<k<1$
$p\in[-1,1)-\{0\}$	$(A_{1,1}+2A)^2_{0,1,0}$, $(A_{1,1}+2A)^2_{\epsilon,1,0}$, $(A_{1,1}+2A)^2_{0,1,\epsilon}$, $(A_{1,1}+2A)^2_{\epsilon,k,-\epsilon}$	$k\in\Re$
C^3	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{\epsilon,0,\epsilon}$	
	$(A_{1,1}+2A)^2_{0,\epsilon,0}$, $(A_{1,1}+2A)^2_{\epsilon,0,-\epsilon}$	
C^4	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{k,0,1}$, $(A_{1,1}+2A)^1_{s,0,-1}$	$0<k,\quad s<0$
	$(A_{1,1}+2A)^2_{0,\epsilon,0}$, $(A_{1,1}+2A)^2_{k,0,1}$, $(A_{1,1}+2A)^2_{s,0,-1}$	$k<0,\quad 0<s$
C_p^5	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{k,0,1}$, $(A_{1,1}+2A)^1_{s,0,-1}$	$0<k,\quad s<0$
$p\geq 0$	$(A_{1,1}+2A)^2_{k,0,1}$, $(A_{1,1}+2A)^2_{s,0,-1}$	$k<0,\quad 0<s$
$(A_{1,1}+2A)^1$	$I_{(1,2)}$	
$(A_{1,1}+2A)^2$	$I_{(1,2)}$	

Where in the above table $\epsilon=\pm 1$.

For three dimensional dual Lie superalgebras $(A_{1,1}+2A)^1_{\alpha,\beta,\gamma}$, $(A_{1,1}+2A)^2_{\alpha,\beta,\gamma}$ where are isomorphic with $(A_{1,1}+2A)^1$, $(A_{1,1}+2A)^2$ respectively, we have the following commutation relations:

$$\{\tilde{X}^2,\tilde{X}^2\}=i\alpha\tilde{X}^1,\quad \{\tilde{X}^2,\tilde{X}^3\}=i\beta\tilde{X}^1,\quad \{\tilde{X}^3,\tilde{X}^3\}=i\gamma\tilde{X}^1,\qquad \alpha,\beta,\gamma\in\Re. \tag{27}$$

Note that these Lie superalgebras are non isomorphic and they differ in the bound of their parameters.

Meanwhile note that for every Lie superalgebra of tables 1 and 2 there are trivial solutions for equations (14) and (15) with $\tilde{f}^{ij}{}_{k}=0$. (i.e. the dual Lie superalgebra is Abelian) we denote these dual Lie superalgebras with $I_{(m,n)}$.

We see that for two and three dimensional Lie super-bialgebras (with two fermions), trivial Lie superalgebras are only dual to nontrivial one. The solution of super Jacobi and mixed super Jacobi identities and related isomorphism matrices C (in step 1) are listed in Appendix B. Here for explanation of steps 1-3 of section 3 we give an example.

5.1 An example

What follows is an explanation of the details of calculations for obtaining dual Lie superalgebras of Lie super-bialgebras $(C^4, (A_{1,1}+2A)^1_{k,0,1})$ and $(C^4, (A_{1,1}+2A)^1_{s,0,-1})$. As mentioned in Appendix B, the solution of super Jacobi and mixed super Jacobi identities have the following form

$$\tilde{f}^{22}{}_1 = i\alpha, \qquad \tilde{f}^{33}{}_1 = i\beta, \qquad \tilde{f}^{23}{}_1 = i\gamma, \quad \forall \alpha, \beta, \gamma \in \Re. \tag{28}$$

This solution is isomorphic with the Lie superalgebra $(A_{1,1}+2A)^1$ with the following matrix:

$$C_1 = \begin{pmatrix} c_{11} & 0 & 0 \\ c_{21} & c_{22} & c_{23} \\ c_{31} & 0 & c_{33} \end{pmatrix}, \tag{29}$$

where $c_{11} = -ic_{33}^2 \tilde{f}^{33}{}_1$, $c_{23} = -c_{22}\frac{\tilde{f}^{23}{}_1}{\tilde{f}^{33}{}_1}$, $\tilde{f}^{22}{}_1 = (\frac{c_{23}^2+c_{33}^2}{c_{22}^2})\tilde{f}^{33}{}_1$ and c_{22}, $c_{33} \in \Re - \{0\}$. By imposing that C_1 must be the transformation matrix (38), we have $c_{31} = c_{21} = 0$.
Now with the help of automorphism supergroup of C^4, the solution of (21) for the matrix B will be

$$B = \begin{pmatrix} \frac{c^2(b_{32}^2+b_{33}^2)}{\beta} & 0 & 0 \\ 0 & \mp\frac{b_{33}c_{33}}{c_{22}} & \frac{b_{33}c_{33}}{c_{22}} \\ 0 & b_{32} & b_{33} \end{pmatrix}, \quad \beta = -i\tilde{f}^{33}{}_1. \tag{30}$$

Now using (20), we obtain the following commutation relations for the dual Lie superalgebra $(A_{1,1}+2A)^{1\prime}$

$$\{\tilde{X}^2, \tilde{X}^2\} = ia^{'}\tilde{X}^1, \qquad \{\tilde{X}^3, \tilde{X}^3\} = ib^{'}\tilde{X}^1, \qquad a^{'} = \frac{\beta c_{33}^2}{c^2 c_{22}^2}, \;\; b^{'} = \frac{\beta}{c^2}, \tag{31}$$

such that $a^{'}$ and $b^{'}$ have the same sings.
One cannot choose $a^{'} = b^{'}$ because in this case we have

$$B_1 = \begin{pmatrix} \frac{c^{'2}(b^{'2}_{32}+b^{'2}_{33})}{\beta} & 0 & 0 \\ 0 & -b^{'}_{33} & b^{'}_{32} \\ 0 & b^{'}_{32} & b^{'}_{33} \end{pmatrix}, \tag{32}$$

so as $BB_1^{-1} \notin A^{st}(C^4)$. Now by choosing $b^{'} = 1$ i.e. $\beta = c^2$ we have

$$B_2 = \begin{pmatrix} b^{''2}_{32} + b^{''2}_{33} & 0 & 0 \\ 0 & \mp\frac{c^{''}_{33}}{c^{'}_{22}}b^{''}_{33} & \frac{c^{''}_{33}}{c^{''}_{22}}b^{''}_{32} \\ 0 & b^{''}_{32} & b^{''}_{33} \end{pmatrix}. \tag{33}$$

such that $BB_2^{-1} \in A^{st}(C^4)$, this means that we can choose $b^{'} = 1$ and in this case we have the following dual Lie superalgebras ($a^{'} = \frac{c_{33}^2}{c_{22}^2} = k > 0$):

$$(A_{1,1}+2A)^1_{k,0,1}: \qquad \{\tilde{X}^2, \tilde{X}^2\} = ik\tilde{X}^1, \qquad \{\tilde{X}^3, \tilde{X}^3\} = i\tilde{X}^1, \;\; k > 0 \tag{34}$$

In the same way by choosing $b^{'} = -1$ i.e. $\beta = -c^2$ such that $a^{'} \neq b^{'}$ we have

$$B_3 = \begin{pmatrix} -(b^{'''2}_{\ \ 32} + b^{'''2}_{\ \ 33}) & 0 & 0 \\ 0 & \mp\frac{c^{'''}_{33}}{c^{'''}_{22}} b^{'''}_{33} & \frac{c^{'''}_{33}}{c^{'''}_{22}} b^{'''}_{32} \\ 0 & b^{'''}_{32} & b^{'''}_{33} \end{pmatrix}. \tag{35}$$

so as $BB_3^{-1} \in A^{st}(C^4)$, this means that we can choose $b^{'} = -1$ and in this case we have the following dual Lie superalgebras ($a^{'} = -\frac{c^2_{33}}{c^2_{22}} = s < 0$):

$$(A_{1,1} + 2A)^1_{s,0,-1} : \qquad \{\tilde{X}^2, \tilde{X}^2\} = is\tilde{X}^1, \qquad \{\tilde{X}^3, \tilde{X}^3\} = -i\tilde{X}^1, \ \ s < 0 \tag{36}$$

Note that as $B_2 B_3^{-1} \notin A^{st}(C^4)$, this means that the Lie super-bialgebras $(C^4, (A_{1,1} + 2A)^1_{k,0,1})$ and $(C^4, (A_{1,1} + 2A)^1_{s,0,-1})$ are not equivalent.

6 Conclusion

We presented a new method for obtaining and classifying low dimensional *Lie super-bialgebras* which can also be applied for classification of low dimensional *Lie bialgebras.* We classified all 48 two and three dimensional Lie super-bialgebras [11]. Using this classification one can construct Poisson-Lie T-dual sigma models on the low dimensional Lie supergroups [19] (as [4] for Lie groups). Moreover, one can determinate the coboundary type of these Lie super-bialgebras and their classical r-matrices and Poisson brackets for their Poisson-Lie supergroups [20]. Determination of doubles and their isomorphism and investigation of Poisson-Lie plurality [21] are open problems for further investigation.

Acknowledgments

We would like to thank S. Moghadassi for carefully reading the manuscript and useful comments.

Appendix A

In this appendix, we will review some basic properties of tensors and matrices from[16].

1. We consider the standard bases for the supervector spaces so that in writing the bases as a column matrix, we first present the bosonic base, then the fermionic one. The transformation of standard bases and its dual bases can be written as follows:

$$e'_i = (-1)^j K_i{}^j e_j, \qquad e'^i = K^{-t\,i}{}_j \, e^j, \tag{37}$$

where the transformation matrix K has the following block diagonal representation [16]

$$K = \left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array}\right), \tag{38}$$

where A, B, C are real submatrices and D is pure imaginary submatrix [16]. Here we consider the matrix and tensors having a form with all upper and lower indices written in the right hand side.

2. The transformation properties of upper and lower right indices to the left one for general tensors are as follows:

$${}^iT^{k}_{jl\ldots} = T^{ik}_{jl\ldots}, \qquad {}_jT^{ik}_{l\ldots} = (-1)^j \, T^{ik}_{jl\ldots}. \tag{39}$$

3. For supertransposition we have

$$L^{t\,i}{}_j = (-1)^{ij} \, L_j{}^i, \qquad L^{t}_i{}^j = (-1)^{ij} \, L^i{}_j,$$

---

[11] Note that in this classification one can omit the $i = \sqrt{-1}$ from commutation relations and to obtain Lie super-bialgebras in the nonstandard bases.

$$M^t_{\ ij} = (-1)^{ij}\ M_{\ ji}, \qquad M^{t\ ij} = (-1)^{ij}\ M^{\ ji}. \tag{40}$$

4. For superdeterminant we have

$$sdet \left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array} \right) = det(A - CB^{-1}D)(detB)^{-1}, \tag{41}$$

when $detB \neq 0$ and

$$sdet \left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array} \right) = (det(B - DA^{-1}C))^{-1}\ (detA), \tag{42}$$

when $detA \neq 0$. For the inverse of matrix we have

$$\left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array} \right)^{-1} = \left(\begin{array}{c|c} (1_m - A^{-1}CB^{-1}D)^{-1}A^{-1} & -(1_m - A^{-1}CB^{-1}D)^{-1}A^{-1}CB^{-1} \\ \hline -(1_n - B^{-1}DA^{-1}C)^{-1}B^{-1}DA^{-1} & (1_n - B^{-1}DA^{-1}C)^{-1}B^{-1} \end{array} \right), \tag{43}$$

where $detA, detB \neq 0$ and m,n are dimensions of submatrices A and B respectively.

Appendix B

This appendix includes solutions of super Jacobi and mixed super Jacobi identities $(sJ - msJ)$ for dual Lie superalgebras and isomorphism matrices C which relate these solutions to other Lie superalgebras.

1. Solutions of $(sJ - msJ)$ for dual Lie superalgebras of B is

$$\tilde{f}^{22}_{\ \ 1} = i\alpha, \qquad \tilde{f}^{12}_{\ \ 1} = 0, \tag{44}$$

where $\alpha \in \Re$.
Isomorphism matrix between this solutions with $(A_{1,1} + A)$ is as follows:

$$C = \left(\begin{array}{cc} -ic_{22}^2 \tilde{f}^{22}_{\ \ 1} & 0 \\ c_{21} & c_{22} \end{array} \right), \tag{45}$$

with the conditions $c_{22} \in \Re - \{0\}$ and imposing that C must be the transformation matrix, we have $c_{21} = 0$.

3. Solutions of $(sJ - msJ)$ for dual Lie superalgebras of C^1_p $(p \in \Re - \{0\})$ are

$i)$ $\quad \tilde{f}^{12}_{\ \ 1} = \alpha, \ \ \tilde{f}^{23}_{\ \ 3} = p\alpha, \qquad p \in \Re - \{0\}, \quad \alpha \in \Re,$

$ii)$ $\quad \tilde{f}^{12}_{\ \ 1} = \tilde{f}^{23}_{\ \ 3} = \beta, \qquad p = 1, \quad \beta \in \Re,$

$iii)$ $\quad \tilde{f}^{33}_{\ \ 1} = i\gamma, \qquad p \in \Re - \{0\}, \quad \gamma \in \Re,$

$iv)$ $\quad \tilde{f}^{33}_{\ \ 1} = i\lambda, \ \ \tilde{f}^{33}_{\ \ 2} = i\eta \qquad p = \frac{1}{2}, \quad \lambda, \eta \in \Re,$

$v)$ $\quad \tilde{f}^{12}_{\ \ 1} = \mu, \ \ \tilde{f}^{23}_{\ \ 3} = -\frac{\mu}{2}, \ \ \tilde{f}^{33}_{\ \ 1} = i\nu \qquad p = -\frac{1}{2}, \quad \mu, \nu \in \Re.$

For solution (i) we have isomorphism matrix between these solutions with C^1_p as follows:

$$C = \left(\begin{array}{ccc} c_{11} & -\frac{1}{\tilde{f}^{12}_{\ \ 1}} & c_{13} \\ c_{21} & 0 & 0 \\ 0 & 0 & c_{33} \end{array} \right), \tag{46}$$

with the conditions $\tilde{f}^{23}_{\ \ 3} = p\tilde{f}^{12}_{\ \ 1}$ and $c_{21}, c_{33} \in \Re - \{0\}$, $c_{11} \in \Re$, as well as imposing that C must be transformation matrix we have $c_{13} = 0$. Solution (ii) is a special case of solution (i). For solution (iii) and (iv) $sdetC = 0$. Solution (v) is investigated for $(C^1_{\frac{1}{2}}, C^1_p)$ in the following step.

3. Solutions of $(sJ - msJ)$ for dual Lie superalgebras of $C^1_{\frac{1}{2}}$ are

$i)$ $\quad \tilde{f}^{23}_{\ \ 3} = \alpha, \ \ \tilde{f}^{12}_{\ \ 1} = 2\alpha, \quad \alpha \in \Re,$

$ii)$ $\quad \tilde{f}^{33}_{\ \ 1} = i\beta, \ \ \tilde{f}^{33}_{\ \ 2} = i\gamma, \ \ \tilde{f}^{23}_{\ \ 3} = \frac{\beta}{2}, \ \ \tilde{f}^{13}_{\ \ 3} = \frac{-\gamma}{2}, \ \ \tilde{f}^{12}_{\ \ 1} = -\beta, \ \ \tilde{f}^{12}_{\ \ 2} = -\gamma, \quad \beta, \gamma \in \Re.$

For solution (i) we have isomorphism matrix C between $(C^1_{\frac{1}{2}}, C^1_p)$ as follows:

$$C = \left(\begin{array}{ccc} c_{11} & -\frac{1}{2\tilde{f}^{23}{}_3} & c_{13} \\ c_{21} & 0 & 0 \\ 0 & 0 & c_{33} \end{array}\right), \tag{47}$$

with the conditions $\tilde{f}^{12}{}_3 = 2\tilde{f}^{23}{}_3$, c_{21}, $c_{33} \in \Re - \{0\}$, $c_{11} \in \Re$ and $p = -\frac{1}{2}$, as well as imposing that C must be the transformation matrix, we have $c_{13} = 0$.

For solution (ii) we have isomorphism matrices C_1 and C_2 between $(C^1_{\frac{1}{2}}, C^1_{\frac{1}{2}})$ as follows:

$$C_1 = \left(\begin{array}{ccc} c_{11} & \frac{i}{\tilde{f}^{33}{}_1} & 0 \\ -ic^2_{33}\tilde{f}^{33}{}_1 & 0 & 0 \\ 0 & 0 & c_{33} \end{array}\right), \tag{48}$$

with the conditions $\tilde{f}^{33}{}_2 = \tilde{f}^{13}{}_3 = \tilde{f}^{12}{}_2 = 0$, $c_{33} \in \Re - \{0\}$ and $c_{11} \in \Re$

$$C_2 = \left(\begin{array}{ccc} c_{12}\frac{\tilde{f}^{33}{}_1}{\tilde{f}^{33}{}_2} - \frac{i}{\tilde{f}^{33}{}_2} & c_{12} & 0 \\ -ic^2_{33}\tilde{f}^{33}{}_1 & -ic^2_{33}\tilde{f}^{33}{}_2 & 0 \\ 0 & 0 & c_{33} \end{array}\right), \tag{49}$$

with the conditions $c_{33} \in \Re - \{0\}$ and $c_{12} \in \Re$.

4. Solution of $(sJ - msJ)$ for dual Lie superalgebras of C^2_p, C^3, C^4 and C^5_p is

$$\tilde{f}^{22}{}_1 = i\alpha, \qquad \tilde{f}^{33}{}_1 = i\beta, \qquad \tilde{f}^{23}{}_1 = i\gamma, \tag{50}$$

where α,β and $\gamma \in \Re$. Furthermore for C^3 we have another solution as follows:

$$\tilde{f}^{22}{}_1 = i\alpha,$$

where for this solution we have $sdetC = 0$ and $\alpha \in \Re$ therefore we omit it.

4.1. For the above solution, we have the following isomorphism matrices C_1, C_2 and C_3 between C^2_p, C^3, C^4 and C^5_p with $(A_{1,1} + 2A)^1$

$$C_1 = \left(\begin{array}{ccc} -ic^2_{33}\tilde{f}^{33}{}_1 & 0 & 0 \\ c_{21} & c_{22} & -c_{22}\frac{\tilde{f}^{23}{}_1}{\tilde{f}^{33}{}_1} \\ c_{31} & 0 & c_{33} \end{array}\right), \tag{51}$$

with the conditions $\tilde{f}^{22}{}_1 = (\frac{c^2_{23}+c^2_{33}}{c^2_{22}})\tilde{f}^{33}{}_1$ and c_{22}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_1 must be the transformation matrix, we have $c_{31} = c_{21} = 0$;

$$C_2 = \left(\begin{array}{ccc} -ic^2_{23}\tilde{f}^{33}{}_1 & 0 & 0 \\ c_{21} & 0 & c_{23} \\ c_{31} & -c_{33}\frac{\tilde{f}^{33}{}_1}{\tilde{f}^{23}{}_1} & c_{33} \end{array}\right), \tag{52}$$

with the conditions $\tilde{f}^{22}{}_1 = (\frac{c^2_{23}+c^2_{33}}{c^2_{32}})\tilde{f}^{33}{}_1$ and c_{23}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_2 must be transformation matrix, we have $c_{31} = c_{21} = 0$;

$$C_3 = \left(\begin{array}{ccc} c_{11} & 0 & 0 \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{array}\right), \tag{53}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{23}^2+c_{33}^2}{c_{22}^2+c_{32}^2})\tilde{f}^{33}{}_{1}$, $\tilde{f}^{23}{}_{1} = -(\frac{c_{33}c_{32}+c_{22}c_{23}}{c_{22}^2+c_{32}^2})\tilde{f}^{33}{}_{1}$ and $c_{11} \in \Re - \{0\}$, c_{22}, c_{23}, c_{32}, $c_{33} \in \Re$ with the conditions $c_{23}^2 + c_{33}^2 \neq 0$, $c_{22}^2 + c_{32}^2 \neq 0$, as well as imposing that C_3 must be the transformation matrix, we have $c_{31} = c_{21} = 0$.

4.2. For the above solution we have isomorphism matrices C_4, C_5, C_6, C_7 and C_8 between C_p^2, C^3, C^4 and C_p^5 with $(A_{1,1} + 2A)^2$ as follows:

$$C_4 = \begin{pmatrix} -ic_{32}(c_{23} - c_{33})\tilde{f}^{23}{}_{1} & 0 & 0 \\ c_{21} & c_{32} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{pmatrix}, \tag{54}$$

with the conditions $\tilde{f}^{22}{}_{1} = -(\frac{c_{23}+c_{33}}{c_{32}})\tilde{f}^{23}{}_{1}$, $\tilde{f}^{33}{}_{1} = 0$, $c_{23} \neq c_{33}$, $c_{32} \in \Re - \{0\}$ and $c_{23}, c_{33} \in \Re$, as well as imposing that C_4 must be the transformation matrix, we have $c_{31} = c_{21} = 0$;

$$C_5 = \begin{pmatrix} ic_{32}(c_{23} + c_{33})\tilde{f}^{23}{}_{1} & 0 & 0 \\ c_{21} & -c_{32} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{pmatrix}, \tag{55}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{23}-c_{33}}{c_{32}})\tilde{f}^{23}{}_{1}$, $\tilde{f}^{33}{}_{1} = 0$, $c_{23} \neq -c_{33}$, $c_{32} \in \Re - \{0\}$ and $c_{23}, c_{33} \in \Re$, as well as imposing that C_5 must be the transformation matrix, we have $c_{31} = c_{21} = 0$;

$$C_6 = \begin{pmatrix} -ic_{23}^2\tilde{f}^{33}{}_{1} & 0 & 0 \\ c_{21} & 0 & c_{23} \\ c_{31} & -c_{33}\frac{\tilde{f}^{33}{}_{1}}{\tilde{f}^{23}{}_{1}} & c_{33} \end{pmatrix}, \tag{56}$$

with the conditions $\tilde{f}^{22}{}_{1} = -(\frac{c_{23}^2-c_{33}^2}{c_{32}^2})\tilde{f}^{33}{}_{1}$ and c_{23}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_6 must be the transformation matrix, we have $c_{31} = c_{21} = 0$;

$$C_7 = \begin{pmatrix} ic_{33}^2\tilde{f}^{33}{}_{1} & 0 & 0 \\ c_{21} & c_{22} & -c_{22}\frac{\tilde{f}^{23}{}_{1}}{\tilde{f}^{33}{}_{1}} \\ c_{31} & 0 & c_{33} \end{pmatrix}, \tag{57}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{23}^2-c_{33}^2}{c_{22}^2})\tilde{f}^{33}{}_{1}$ and c_{22}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_7 must be the transformation matrix, we have $c_{31} = c_{21} = 0$;

$$C_8 = \begin{pmatrix} c_{11} & 0 & 0 \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{pmatrix}, \tag{58}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{33}^2-c_{23}^2}{c_{32}^2-c_{22}^2})\tilde{f}^{33}{}_{1}$, $\tilde{f}^{23}{}_{1} = (\frac{c_{22}c_{23}-c_{32}c_{33}}{c_{32}^2-c_{22}^2})\tilde{f}^{33}{}_{1}$ and $c_{11} \in \Re - \{0\}$, c_{22}, c_{23}, c_{32}, $c_{33} \in \Re$ with the conditions $c_{33}^2 - c_{23}^2 \neq 0, c_{32}^2 - c_{22}^2 \neq 0$, as well as imposing that C_8 must be the transformation matrix, we have $c_{31} = c_{21} = 0$.

References

[1] V. G. Drinfel'd, *Hamiltonian Lie groups, Lie bialgebras and the geometric meaning of the classical Yang-Baxter equation*, Sov. Math. Dokl. **27** (1983) 68-71.

[2] P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras,* I, Selecta Math. **2** (1996) n. 1, 1-41.
P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras*, II. Selecta Math. **4** (1998) no. 2, 213-231.

[3] P. Delorme, *Sur les triples de Manin pour les complexes,* Jour. Alg. **246** (2001) 97-174.

[4] M.A. Jafarizadeh and A. Rezaei-Aghdam, *Poisson-Lie T-duality and Bianchi type algebras*, *Phys. Lett. B* **458** (1999) 477-490, `[arXiv:hep-th/9903152]`.

[5] X. Gomez, *Classification of three-dimensional Lie bialgebras*, J. Math. Phys. **41** (2000) 493956.

[6] L. Hlavaty and L. Snobl, *Classification of 6-dimensional Manin triples*, `[arXiv:math-ph/0202209]`.

[7] A. Rezaei-Aghdam, M. Hemmati, A. R. Rastkar, *classification of real three-dimensional Lie bialgebras and their Poisson-Lie groups*, *J. Phys. A:Math.Gen.* **38** (2005) 3981-3994, `[arXiv:math-ph/0412092]`.

[8] Y. Kosmann-Schwarzbach, *Lie bialgebras, Poisson-Lie groups and dressing transformations integrability of nonlinear Systems*: Proc.(Pondicherry) ed Y Kosmann-Schwarzbach, B Grammaticos and K M Tamizhmani (Berlin: Springer)pp 104-70.

[9] N. Andruskiewitsch, *Lie Superbialgebras and Poisson-Lie supergroups*, Abh. Math. Sem. Univ. Hamburg **63** (1993) 147-163.

[10] N. Beisert and E. Spill, *The classical r-matrices of AdS/CFT and its Lie bialgebras structure*, `[arXiv:0708.1762 [hep-th]]`.

[11] N. Geer, *Etingof-Kazdan quantization of Lie super bialgebras*, `[arXiv:math.QA/0409563]`.

[12] C. Juszczak and J. T. Sobczyk, *Classification of low dimentional Lie super-bialgebras, J. Math. Phys.* **39**, (1998) 4982-4992.

[13] C. Juszczak, *Classification of* $osp(2|2)$ *Lie super-bialgebras,* `[arXiv:math.QA/9906101]`.

[14] G. Karaali, *A New Lie Bialgebra Structure on sl(2,1), Contemp. Math.* **413** (2006), pp.101-122.

[15] N. Backhouse, *A classification of four-dimensional Lie superalgebras*, *J. Math. Phys.* **19** (1978) 2400-2402.

[16] B. DeWitt, *Supermanifold,* Cambridge University Press 1992.

[17] A. Eghbali and A. Rezaei-Aghdam, *Classification of four-dimensional Lie super-bialgebras of type* $(2, 2)$, work in progress.

[18] C. Klimčik and P. Ševera, *Dual non-Abelian duality and the Drinfeld double*, *Phys. Lett.* **B351** (1995) 445-462, `[arXiv:hep-th/9502122]`.
C. Klimčik, *Poisson-Lie T-duality*, *Nucl. Phys. Proc. Suppl.* **46** (1996) 116-121, `[arXiv:hep-th/9509095]`.

[19] A. Eghbali, A. Rezaei-Aghdam, *Poisson-Lie T-dual sigma models on supermanifolds*, JHEP 0909:094 (2009), `[arXiv:0901.1592v3 [hep-th]]`.

[20] A. Eghbali and A. Rezaei-Aghdam, *Classical r-matrices of two and three dimensional Lie super-bialgebras and their Poisson-Lie supergroups*, `[ arXiv:0908.2182 [math-ph]]`.

[21] R. Von Unge, *Poisson-Lie T-plurality*, JHEP **07**(2002)014 (preprint `[arXiv:hep-th/0205245]`).

arXiv:0901.4471v4 [math-ph] 19 Jul 2010

Classification of two and three dimensional Lie super-bialgebras

A. Eghbali [*], A. Rezaei-Aghdam [†] and F. Heidarpour

Department of Physics, Faculty of science, Azarbaijan University
of Tarbiat Moallem , 53714-161, Tabriz, Iran

July 20, 2010

Abstract

Using adjoint representation of Lie superalgebras, we obtain the matrix form of super Jacobi and mixed super Jacobi identities of Lie super-bialgebras. By direct calculations of these identities, and use of automorphism supergroups of two and three dimensional Lie superalgebras, we obtain and classify all two and three dimensional Lie super-bialgebras.

---

[*] e-mail: a.eghbali@azaruniv.edu

[†] Corresponding author. e-mail: rezaei-a@azaruniv.edu

1 Introduction

Lie bialgebras were first introduced, from the mathematical point of view, by Drinfel'd as algebraic structures and classical limit of underlying quantized enveloping algebras (*quantum groups*) [1]. In particular, every deformation of a universal enveloping algebra induces a Lie bialgebra structure on the underlying Lie algebra. Conversely, it has been shown in Ref. [2] that each Lie bialgebra admits quantization. So the classification of Lie bialgebras can be seen as the first step in the classification of quantum groups. Many interesting examples of Lie bialgebras based on complex semisimple Lie algebras have been given in Drinfel'd [1]. A complete classification of Lie bialgebras with reduction was given in Ref. [3]. However, a classification of Lie bialgebras is out of reach, with similar reasons as for Lie algebra classification. In the non-semisimple case, only a bunch of low dimensional examples have been thoroughly studied [4, 5, 6, 7]. On the other hand, from the physical point of view, the theory of classical integrable systems naturally relates to the geometry and representation theory of Poisson-Lie groups and the corresponding Lie bialgebras and their classical r-matrices (see, for example, Ref. [8]). In the same way, Lie super-bialgebras [9], as the underlying symmetry algebras, play an important role in the integrable structure of AdS/CFT correspondence [10]. In this way, and by considering that there is a universal quantization for Lie super-bialgebras [11], one can assign an important role to the classification of Lie super-bialgebras (especially low dimensional Lie super-bialgebras) from both physical and mathematical point of view. There are distinguished and nonsystematic ways for obtaining low dimensional Lie super-bialgebras (see, for example, Refs. [12, 13, 14]). In this paper, using the adjoint representation of Lie superalgebras, we present a systematic way for obtaining and classifying low dimensional Lie super-bialgebras. We apply this method to the classification of two and three dimensional Lie super-bialgebras.

The paper is organized as follows. In section two, we give basic definitions and notations that are used throughout the paper. The systematic way for classification of Lie super-bialgebras by using matrix form of super Jacobi (sJ) and mixed super Jacobi (msJ) identities of Lie super-bialgebras is described in section three. A list of two and three dimensional Lie superalgebras of Ref. [15] is offered in section 4. The automorphism Lie supergroups of these Lie superalgebras are also presented in section 4 . Then, using the method mentioned in section two, we classify all 48 two and three dimensional Lie super-bialgebras in section five. The details of calculations are explained, using an example. The concluding section discusses some remarks. Certain properties of tensors and supermatrices are given in appendix A. Solutions of ($sJ - msJ$) identities and their isomorphism matrices are give in appendix B.

2 Basic definitions and notations

In this paper we use DeWitt notation for supervector spaces, supermatrices, etc [16]. Some of definitions and related notations are given in appendix A.

Definition: A *Lie superalgebra* $\mathbf{g}$ is a graded vector space $\mathbf{g} = \mathbf{g}_B \oplus \mathbf{g}_F$ with gradings; $grade(\mathbf{g}_B) = 0$, $grade(\mathbf{g}_F) = 1$; such that Lie bracket satisfies the super antisymmetric and super Jacobi identities, i.e., in a graded basis $\{X_i\}$ of $\mathbf{g}$ if we put [17]

$$[X_i, X_j] = f^k{}_{ij} X_k, \tag{1}$$

then

$$(-1)^{i(j+k)} f^m{}_{jl} f^l{}_{ki} + f^m{}_{il} f^l{}_{jk} + (-1)^{k(i+j)} f^m{}_{kl} f^l{}_{ij} = 0, \tag{2}$$

so that

$$f^k{}_{ij} = -(-1)^{ij} f^k{}_{ji}. \tag{3}$$

Note that, in the standard basis, $f^B{}_{BB}$ and $f^F{}_{BF}$ are real c-numbers and $f^B{}_{FF}$ are pure imaginary c-numbers and other components of structure constants $f^i{}_{jk}$ are zero [16], i.e. we have

$$f^k{}_{ij} = 0, \qquad if \quad grade(i) + grade(j) \neq grade(k) \quad (mod 2). \tag{4}$$

Let $\mathbf{g}$ be a finite-dimensional Lie superalgebra and $\mathbf{g}^*$ be its dual vector space and let $(. \ , \ .)$ be the canonical pairing on $\mathbf{g}^* \oplus \mathbf{g}$.

Definition: A *Lie super-bialgebra* structure on a Lie superalgebra $\mathbf{g}$ is a super skew-symmetric linear map $\delta : \mathbf{g} \longrightarrow \mathbf{g} \otimes \mathbf{g}$ (*the super cocommutator*) so that [9]

1) δ is a super one-cocycle, i.e.

$$\delta([X,Y]) = (ad_X \otimes I + I \otimes ad_X)\delta(Y) - (-1)^{|X||Y|}(ad_Y \otimes I + I \otimes ad_Y)\delta(X) \qquad \forall X, Y \in \mathbf{g}, \tag{5}$$

where $|X|(|Y|)$ indicates the grading of $X(Y)$;

2) the dual map ${}^t\delta : \mathbf{g}^* \otimes \mathbf{g}^* \to \mathbf{g}^*$ is a Lie superbracket on $\mathbf{g}^*$, i.e.,

$$(\xi \otimes \eta, \delta(X)) = ({}^t\delta(\xi \otimes \eta), X) = ([\xi, \eta]_*, X) \qquad \forall X \in \mathbf{g};\ \ \xi, \eta \in \mathbf{g}^*. \tag{6}$$

The Lie super-bialgebra defined in this way will be indicated by $(\mathbf{g}, \mathbf{g}^*)$ or $(\mathbf{g}, \delta)$.

Proposition: If there exists an automorphism A of $\mathbf{g}$ such that

$$\delta' = (A \otimes A) \circ \delta \circ A^{-1}, \tag{7}$$

then the super one-cocycles δ and δ' of the Lie superalgebra $\mathbf{g}$ are *equivalent* [18]. In this case the two Lie super-bialgebras $(\mathbf{g}, \delta)$ and $(\mathbf{g}, \delta')$ are equivalent (as in the bosonic case [7]).

Definition: A *Manin* super triple [9] is a triple of Lie superalgebras $(\mathcal{D}, \mathbf{g}, \tilde{\mathbf{g}})$ together with a non-degenerate ad-invariant super symmetric bilinear form $< . \, , \, . >$ on $\mathcal{D}$ such that

1) $\mathbf{g}$ and $\tilde{\mathbf{g}}$ are Lie sub-superalgebras of $\mathcal{D}$,

2) $\mathcal{D} = \mathbf{g} \oplus \tilde{\mathbf{g}}$ as a supervector space,

3) $\mathbf{g}$ and $\tilde{\mathbf{g}}$ are isotropic with respect to $< . \, , \, . >$, i.e.

$$< X_i, X_j > = < \tilde{X}^i, \tilde{X}^j > = 0, \qquad {\delta_i}^j = < X_i, \tilde{X}^j > = (-1)^{ij} < \tilde{X}^j, X_i > = (-1)^{ij} {\delta^j}_i, \tag{8}$$

where $\{X_i\}$ and $\{\tilde{X}^i\}$ are basis of Lie superalgebras $\mathbf{g}$ and $\tilde{\mathbf{g}}$, respectively. Note that in the above relation ${\delta^j}_i$ is the ordinary delta function. There is a one-to-one correspondence between Lie super-bialgebra $(\mathbf{g}, \mathbf{g}^*)$ and Manin super triple $(\mathcal{D}, \mathbf{g}, \tilde{\mathbf{g}})$ with $\mathbf{g}^* = \tilde{\mathbf{g}}$ [9]. If we choose the structure constants of Lie superalgebras $\mathbf{g}$ and $\tilde{\mathbf{g}}$ as

$$[X_i, X_j] = {f^k}_{ij} X_k, \qquad\qquad [\tilde{X}^i, \tilde{X}^j] = {\tilde{f}^{ij}}_{\ \ k} \tilde{X}^k, \tag{9}$$

then ad-invariance of the bilinear form $< . \, , \, . >$ on $\mathcal{D} = \mathbf{g} \oplus \tilde{\mathbf{g}}$ implies that

$$[X_i, \tilde{X}^j] = (-1)^j {\tilde{f}^{jk}}_{\ \ i} X_k + (-1)^i {f^j}_{ki} \tilde{X}^k. \tag{10}$$

Clearly, using Eqs. (6) and (9) we have

$$\delta(X_i) = (-1)^{jk} {\tilde{f}^{jk}}_{\ \ i} X_j \otimes X_k, \tag{11}$$

note that the appearance of $(-1)^{jk}$ in this relation is due to the definition of natural inner product between $\mathbf{g} \otimes \mathbf{g}$ and $\mathbf{g}^* \otimes \mathbf{g}^*$ as $(\tilde{X}^i \otimes \tilde{X}^j, X_k \otimes X_l) = (-1)^{jk} {\delta^i}_k {\delta^j}_l$.

As a result, if we apply this relation in the super one-cocycle condition (5), super Jacobi identities (2) for the dual Lie superalgabra and the following mixed super Jacobi identities are obtained

$${f^m}_{jk} {\tilde{f}^{il}}_{\ \ m} = {f^i}_{mk} {\tilde{f}^{ml}}_{\ \ \ j} + {f^l}_{jm} {\tilde{f}^{im}}_{\ \ \ k} + (-1)^{jl} {f^i}_{jm} {\tilde{f}^{ml}}_{\ \ \ k} + (-1)^{ik} {f^l}_{mk} {\tilde{f}^{im}}_{\ \ \ j}. \tag{12}$$

This relation can also be obtained from super Jacobi identity of $\mathcal{D}$.

3 Calculation of Lie super-bialgebras using adjoint representation

As discussed in the introduction, by use of super Jacobi and mixed super Jacobi identities we classify two and three dimensional Lie super-bialgebras. This method was first used to obtain three dimensional Lie bialgebras in Ref. [4]. Because of tensorial form of super Jacobi and mixed super Jacobi identities, working with them is not so easy and we suggest writing these equations as matrix forms using the following adjoint representations for Lie superalgebras $\mathbf{g}$ and $\tilde{\mathbf{g}}$

$$(\tilde{\mathcal{X}}^i)^j{}_k = -\tilde{f}^{ij}{}_k, \qquad (\mathcal{Y}^i)\,{}_{jk} = -f^i{}_{jk}. \tag{13}$$

Then the matrix forms of super Jacobi identities (2) for dual Lie superalgebra $\tilde{\mathbf{g}}$ and mixed super Jacobi identities (12) become as follows, respectively:

$$(\tilde{\mathcal{X}}^i)^j{}_k \tilde{\mathcal{X}}^k - \tilde{\mathcal{X}}^j \tilde{\mathcal{X}}^i + (-1)^{ij} \tilde{\mathcal{X}}^i \tilde{\mathcal{X}}^j = 0, \tag{14}$$

$$(\tilde{\mathcal{X}}^i)^j{}_l\, \mathcal{Y}^l = -(-1)^k (\tilde{\mathcal{X}}^{st})^j\, \mathcal{Y}^i + \mathcal{Y}^j \tilde{\mathcal{X}}^i - (-1)^{ij} \mathcal{Y}^i \tilde{\mathcal{X}}^j + (-1)^{k+ij} (\tilde{\mathcal{X}}^{st})^i\, \mathcal{Y}^j. \tag{15}$$

Note that in the above relations we use the right indices for the matrix elements and index k represents the column of matrix $\tilde{\mathcal{X}}^{st}$.

Having the structure constants of the Lie superalgebra $\mathbf{g}$, we solve the matrix Eqs. (14) and (15) in order to obtain the structure constants of the dual Lie superalgebras $\tilde{\mathbf{g}}$ so that $(\mathbf{g}, \tilde{\mathbf{g}})$ is Lie super-bialgebra. We assume that, we have classification and tabulation of the Lie superalgebras $\mathbf{g}$ (for example two and three dimensional Lie superalgebras for this paper). Here we explain the method in general and one can apply it for classification of other low dimensional Lie super-bialgebras [19]. We fulfill this work in the following three steps. Note that in solving Eqs. (14), (15), (17) and (21) we use the MAPLE program.

Step 1: Solutions of super Jacobi and mixed super Jacobi identities and determination of Lie superalgebras $\mathbf{g}'$ *which are isomorphic with dual solutions*

With the solution of matrix Eqs. (14) and (15) for obtaining matrices $\tilde{\mathcal{X}}^i$, some structure constants of $\tilde{\mathbf{g}}$ are obtained to be zero, some unknown and some obtained in terms of each other. In order to know whether $\tilde{\mathbf{g}}$ is one of the Lie superalgebras of table or isomorphic to them, we must use the following isomorphic relation between obtained Lie superalgebras $\tilde{\mathbf{g}}$ and one of the Lie superalgebras of table, e.g., $\mathbf{g}'$. Applying the transformation (37) for a change of basis $\tilde{\mathbf{g}}$ we have

$$\tilde{X}'^{\,i} = C^i{}_j \tilde{X}^j, \qquad [\tilde{X}'^{\,i}, \tilde{X}'^{\,j}] = \tilde{f}'^{\,ij}_{\;k} \tilde{X}'^{\,k}, \tag{16}$$

then the following matrix equations for isomorphism

$$(-1)^{i(j+l)}\, C\, (C^i{}_k\, \tilde{\mathcal{X}}^k_{(\tilde{\mathbf{g}})}) = \mathcal{X}^i_{(\mathbf{g}')}\, C, \tag{17}$$

are obtained, where the indices j and l correspond to the row and column of C in the left hand side of (17). In the above matrix equations $\mathcal{X}^i_{(\mathbf{g}')}$ are adjoint matrices of known Lie superalgebra $\mathbf{g}'$ of the classification table. Solving (17) with the condition $sdetC \neq 0$ we obtain some extra conditions on $\tilde{f}^{kl}_{(\tilde{\mathbf{g}})}{}_m$'s that were obtained from (14) and (15).

Step 2: Obtaining general form of the transformation matrices $B : \mathbf{g}' \longrightarrow \mathbf{g}'.i$*; such that* $(\mathbf{g}, \mathbf{g}'.i)$ *are Lie super-bialgebras*

As the second step we transform Lie super-bialgebra $(\mathbf{g}, \tilde{\mathbf{g}})$ (where in the Lie superalgebra $\tilde{\mathbf{g}}$ we impose extra conditions obtained in the step one) to Lie super bialgebra $(\mathbf{g}, \mathbf{g}'.i)$ (where $\mathbf{g}'.i$ is isomorphic as Lie superalgebra to $\mathbf{g}'$) with an automorphism of the Lie superalgebra $\mathbf{g}$. As the inner product (8) is invariant we have $A^{-st} : \tilde{\mathbf{g}} \longrightarrow \mathbf{g}'.i$,

$$X'_i = (-1)^k A_i{}^k X_k, \qquad \tilde{X}'^{\,j} = (A^{-st})^j{}_l \tilde{X}^l, \qquad < X'_i, \tilde{X}'^{\,j} >= \delta_i{}^j, \tag{18}$$

where A^{-st} is superinverse supertranspose of every matrix $A \in Aut(\mathbf{g})$ [$Aut(\mathbf{g})$ is the automorphism supergroup of the Lie superalgebra $\mathbf{g}$]. Thus we have the following transformation relation for the map A^{-st}:

$$(-1)^{k(j+l)} (A^{-st})^i{}_k \tilde{f}^{kl}_{(\tilde{\mathbf{g}})}{}_m (A^{-st})^j{}_l = f^{ij}{}_{(\mathbf{g}'.\mathbf{i})}{}_n (A^{-st})^n{}_m. \tag{19}$$

Now, for obtaining Lie super-bialgebras $(\mathbf{g}, \mathbf{g}'.i)$, we must find Lie superalgebras $\mathbf{g}'.i$ or transformations $B : \mathbf{g}' \longrightarrow \mathbf{g}'.i$, such that

$$(-1)^{k(j+l)} B^i{}_k f^{kl}{}_{(\mathbf{g}')\; m} B^j{}_l = f^{ij}{}_{(\mathbf{g}'.\mathbf{i})\; n} B^n{}_m. \tag{20}$$

For this purpose, it is enough to omit $f^{ij}{}_{(\mathbf{g}'.\mathbf{i})\; n}$ between (19) and (20). Then we will have the following matrix equation for B:

$$(A^{-st})^i{}_m \tilde{\mathcal{X}}^{st\; m}_{(\tilde{\mathbf{g}})} A^{-1} = (-1)^{i(l+j)} (B^{st} A)^{-1} (B^i{}_k \mathcal{X}^{st\; k}{}_{(\mathbf{g}')}) B^{st}, \tag{21}$$

where indices l and j are row and column of B^{st} in the right hand side of the above relation. Note that for A^{-st} and A^{-1} in (21) we use row and column indices on the right hand sides. Now by solving (21) we obtain the general form of matrix B with the condition $sdetB \neq 0$. In solving (21) one can obtain conditions on elements of matrix A, yet we must only consider those conditions under which we have $sdetA \neq 0$ and matrices A, B and A^{-st} have the general transformation matrix form (38).

Step 3: Obtaining and classificating the nonequivalent Lie super-bialgebras

Having solved (21), we obtain the general form of the matrix B so that its elements are written in terms of the elements of matrices A, C and structure constants $\tilde{f}^{ij}_{(\tilde{\mathbf{g}})\; k}$. Now with substituting B in (20), we obtain structure constants $f^{ij}{}_{(\mathbf{g}'.\mathbf{i})\; n}$ of the Lie superalgebra $\mathbf{g}'.i$ in terms of elements of matrices A and C and some $\tilde{f}^{ij}_{(\tilde{\mathbf{g}})\; k}$. Then we check whether it is possible to equalize the structure constants $f^{ij}{}_{(\mathbf{g}'.\mathbf{i})\; n}$ with each other and with ± 1 or not so as to remark $sdetB \neq 0$, $sdetA \neq 0$, and $sdetC \neq 0$. In this way, we obtain isomorphism matrices B_1, B_2,.... As B_i^{st} must be in the form of transformation matrices (38), we obtain conditions on the matrices B_i. The reason is that if $(\mathbf{g}, \mathbf{g}'.i)$ is Lie super-bialgebra, then $(\mathbf{g}', \mathbf{g}.i)$ will be Lie super-bialgebras with $B_i^{st} : \mathbf{g} \longrightarrow \mathbf{g}.i$. Note that in obtaining B_is we impose the condition $B{B_i}^{-1} \in Aut^{st}(\mathbf{g})$ [$Aut^{st}(\mathbf{g})$ is the supertranspose of $Aut(\mathbf{g})$]; if this condition is not satisfied then we cannot impose it on the structure constants because B and B_i are not equivalent (see bellow).

Now using isomorphism matrices B_1, B_2, etc, we can obtain Lie super-bialgebras $(\mathbf{g}, \mathbf{g}'.i)$, $(\mathbf{g}, \mathbf{g}'.ii)$, etc. On the other hand, there is the question: which of these Lie super-bialgebras are equivalent? In order to answer this question, we use the matrix form of the relation (7). Consider the two Lie super-bialgebras $(\mathbf{g}, \mathbf{g}'.i)$, $(\mathbf{g}, \mathbf{g}'.ii)$; then using [20]

$$A(X_i) = (-1)^j A_i{}^j X_j, \tag{22}$$

relation (7) will have the following matrix form:

$$(-1)^{i(j+l)} A^{st}((A^{st})^i{}_k \mathcal{X}_{(\mathbf{g}'.i)}{}^k) = \mathcal{X}_{(\mathbf{g}'.ii)}{}^i A^{st}. \tag{23}$$

On the other hand, the transformation matrix between $\mathbf{g}'.i$ and $\mathbf{g}'.ii$ is $B_2 B_1^{-1}$ if $B_1 : \mathbf{g}' \longrightarrow \mathbf{g}'.i$ and $B_2 : \mathbf{g}' \longrightarrow \mathbf{g}'.ii$; then we have

$$(-1)^{i(j+l)} (B_2 B_1^{-1})((B_2 B_1^{-1})^i{}_k \mathcal{X}_{(\mathbf{g}'.i)}{}^k) = \mathcal{X}_{(\mathbf{g}'.ii)}{}^i (B_2 B_1^{-1}). \tag{24}$$

A comparison of (24) with (23) reveals that if $B_2 B_1^{-1} \in A^{st}$ holds, then the Lie super-bialgebras $(\mathbf{g}, \mathbf{g}'.i)$ and $(\mathbf{g}, \mathbf{g}'.ii)$ are equivalent. In this way, we obtain nonequivalent class of B_is and we consider only one element of this class. Thus, we obtain and classify all Lie super-bialgebras. In the next section, we apply this formulation to two and three dimensional Lie superalgebras.

4 Two and three dimensional Lie superalgebras and their automorphism supergroups

In this section, we use the classification of two and three dimensional Lie superalgebras listed in Ref. [15]. In this classification, Lie superalgebras are divided into two types: trivial and nontrivial Lie superalgebras for which the commutations of fermion-fermion is zero or nonzero, respectively (as we use DeWitt notation here, the structure constant C^B_{FF} must be pure imaginary). The results have been presented in tables 1 and 2. As the tables $(m, n-m)$ indicate, the Lie superalgebras have $m, \{X_1, ..., X_m\}$ bosonic and $n-m, \{X_{m+1}, ..., X_n\}$ fermionic generators.

Table 1 : Trivial Lie superalgebras.

Type	**g**	Bosonic basis	Fermionic basis	Non-zero commutation relations	Comments
$(1,1)$	B	X_1	X_2	$[X_1, X_2] = X_2$	
$(2,1)$	C_p^1	X_1, X_2	X_3	$[X_1, X_2] = X_2,\ [X_1, X_3] = pX_3$	$p \neq 0$
$(1,2)$	C_p^2	X_1	X_2, X_3	$[X_1, X_2] = X_2,\ [X_1, X_3] = pX_3$	$0 < \lvert p \rvert \leq 1$
	C^3	X_1	X_2, X_3	$[X_1, X_3] = X_2$	Nilpotent
	C^4	X_1	X_2, X_3	$[X_1, X_2] = X_2,\ [X_1, X_3] = X_2 + X_3$	
	C_p^5	X_1	X_2, X_3	$[X_1, X_2] = pX_2 - X_3,\ [X_1, X_3] = X_2 + pX_3$	$p \geq 0$

Table 2 : Nontrivial Lie superalgebras ([21]).

Type	**g**	Bosonic basis	Fermionic basis	Non-zero (anti)commutation relations	Comments
$(1,1)$	$(A_{1,1} + A)$	X_1	X_2	$\{X_2, X_2\} = iX_1$	
$(2,1)$	$C^1_{\frac{1}{2}}$	X_1, X_2	X_3	$[X_1, X_2] = X_2,\ [X_1, X_3] = \frac{1}{2}X_3,\ \{X_3, X_3\} = iX_2$	
$(1,2)$	$(A_{1,1} + 2A)^1$	X_1	X_2, X_3	$\{X_2, X_2\} = iX_1,\ \{X_3, X_3\} = iX_1$	Nilpotent
	$(A_{1,1} + 2A)^2$	X_1	X_2, X_3	$\{X_2, X_2\} = iX_1,\ \{X_3, X_3\} = -iX_1$	Nilpotent

One can check whether super Jacobi identities for the above Lie superalgebras are satisfied. As mentioned in section 3 for obtaining dual Lie superalgebras we need automorphism supergroups of Lie superalgebras. In order to calculate the automorphism supergroups of two and three dimensional Lie superalgebras, we use the following transformation:

$$X'_i = (-1)^j \, A_i^{\ j} X_j, \qquad\qquad [X'_i, X'_j] = f^k_{\ ij} X'_k, \tag{25}$$

thus we have the following matrix equation for elements of automorphism supergroups:

$$(-1)^{ij+mk} A \mathcal{Y}^k A^{st} = \mathcal{Y}^e A_e^{\ k}, \tag{26}$$

where index j is the column of matrix A and indices i and m are row and column of A^{st} in the left hand side. Using (26) with the condition $sdetA \neq 0$ and imposing the condition that A must be in the form (38), we obtain table 3 for automorphism supergroups.

Table 3 : Automorphism supergroups of the two and three dimensional Lie superalgebras.

$\mathbf{g}$	Automorphism supergroups	Comments
B	$\begin{pmatrix} 1 & 0 \\ 0 & b \end{pmatrix}$	$b \in \Re - \{0\}$
$(A_{1,1}+A)$	$\begin{pmatrix} a^2 & 0 \\ 0 & a \end{pmatrix}$	$a \in \Re - \{0\}$
C_p^1	$\begin{pmatrix} 1 & a & 0 \\ 0 & c & 0 \\ 0 & 0 & d \end{pmatrix}$	$p \in \Re - \{0\}, \quad c, d \in \Re - \{0\}, \; a \in \Re$
$C_{\frac{1}{2}}^1$	$\begin{pmatrix} 1 & a & 0 \\ 0 & b^2 & 0 \\ 0 & 0 & b \end{pmatrix}$	$b \in \Re - \{0\}$, $a \in \Re$
C_1^2	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & b \\ 0 & a & d \end{pmatrix}$	$ab - cd \neq 0$
C_p^2	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & d \end{pmatrix}$	$c, d \in \Re - \{0\}, \; p \in [-1,1] - \{0\}$
C^3	$\begin{pmatrix} a & 0 & 0 \\ 0 & ad & 0 \\ 0 & e & d \end{pmatrix}$	$a, d \in \Re - \{0\}$, $e \in \Re$
C^4	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & 0 \\ 0 & d & c \end{pmatrix}$	$c \in \Re - \{0\}, \; d \in \Re$
C_p^5	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & c & -d \\ 0 & d & c \end{pmatrix}$	$p \geq 0, \; c \in \Re - \{0\}$ or $d \in \Re - \{0\}$
$(A_{1,1}+2A)^1$	$\begin{pmatrix} b^2+c^2 & 0 & 0 \\ 0 & b & -c \\ 0 & c & b \end{pmatrix}$	$b \in \Re - \{0\}$ or $c \in \Re - \{0\}$
$(A_{1,1}+2A)^2$	$\begin{pmatrix} b^2-c^2 & 0 & 0 \\ 0 & b & c \\ 0 & c & b \end{pmatrix}$	$b \in \Re - \{0\}$ or $c \in \Re - \{0\}$

5 Two and three dimensional Lie super-bialgebras

Using the methods discussed in section 3, we can classify two and three dimensional Lie super-bialgebras. We have applied MAPLE 9 and obtained 48 Lie super-bialgebras. These have been listed in the following tables 4-6:

Table 4 : Two dimensional Lie super-bialgebras of the type $(1,1)$.

$\mathbf{g}$	$\tilde{\mathbf{g}}$	Non-zero (anti) commutation relations of $\tilde{\mathbf{g}}$
$(A_{1,1}+A)$	$I_{(1,1)}$	
B	$I_{(1,1)}$	
	$(A_{1,1}+A)$	$\{\tilde{X}^2, \tilde{X}^2\} = i\tilde{X}^1$
	$(A_{1,1}+A).i$	$\{\tilde{X}^2, \tilde{X}^2\} = -i\tilde{X}^1$

Table 5: Three dimensional Lie super-bialgebras of the type (2 ,1).

$\mathbf{g}$	$\tilde{\mathbf{g}}$	Non-zero (anti) commutation relations of $\tilde{\mathbf{g}}$	Comments
C_p^1	$I_{(2,1)}$		
	$C^1_{-p}.i$	$[\tilde{X}^1, \tilde{X}^2] = \tilde{X}^1, \quad [\tilde{X}^2, \tilde{X}^3] = p\tilde{X}^3$	$p \in \Re - \{0\}$
$C^1_{\frac{1}{2}}$	$I_{(2,1)}$		
	$C^1_p.i\vert_{p=-\frac{1}{2}}$	$[\tilde{X}^1, \tilde{X}^2] = \tilde{X}^1, \quad [\tilde{X}^2, \tilde{X}^3] = \frac{1}{2}\tilde{X}^3$	
	$C^1_p.ii\vert_{p=-\frac{1}{2}}$	$[\tilde{X}^1, \tilde{X}^2] = -\tilde{X}^1, \quad [\tilde{X}^2, \tilde{X}^3] = -\frac{1}{2}\tilde{X}^3$	
	$C^1_{\frac{1}{2}}.i$	$[\tilde{X}^1, \tilde{X}^2] = \tilde{X}^1, \quad [\tilde{X}^2, \tilde{X}^3] = -\frac{1}{2}\tilde{X}^3, \quad \{\tilde{X}^3, \tilde{X}^3\} = i\tilde{X}^1$	
	$C^1_{\frac{1}{2}}.ii$	$[\tilde{X}^1, \tilde{X}^2] = -\tilde{X}^1, \quad [\tilde{X}^2, \tilde{X}^3] = \frac{1}{2}\tilde{X}^3, \quad \{\tilde{X}^3, \tilde{X}^3\} = -i\tilde{X}^1$	
	$C^1_{\frac{1}{2},k}$	$[\tilde{X}^1, \tilde{X}^2] = k\tilde{X}^2, \ [\tilde{X}^1, \tilde{X}^3] = \frac{k}{2}\tilde{X}^3, \ \{\tilde{X}^3, \tilde{X}^3\} = ik\tilde{X}^2$	$k \in \Re - \{0\}$

Table 6: Three dimensional Lie super-bialgebras of the type (1 ,2). Where $\epsilon = \pm 1$.

$\mathbf{g}$	$\tilde{\mathbf{g}}$	Comments
C_1^2	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{\epsilon,0,\epsilon}$	
	$(A_{1,1}+2A)^2_{\epsilon,0,-\epsilon}$	
C_p^2 $p \in [-1,1) - \{0\}$	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{\epsilon,k,\epsilon}$	$-1 < k < 1$
	$(A_{1,1}+2A)^2_{0,1,0}$, $(A_{1,1}+2A)^2_{\epsilon,1,0}$, $(A_{1,1}+2A)^2_{0,1,\epsilon}$, $(A_{1,1}+2A)^2_{\epsilon,k,-\epsilon}$	$k \in \Re$
C^3	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{\epsilon,0,\epsilon}$	
	$(A_{1,1}+2A)^2_{0,\epsilon,0}$, $(A_{1,1}+2A)^2_{\epsilon,0,-\epsilon}$	
C^4	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{k,0,1}$, $(A_{1,1}+2A)^1_{s,0,-1}$	$0 < k, \quad s < 0$
	$(A_{1,1}+2A)^2_{0,\epsilon,0}$, $(A_{1,1}+2A)^2_{k,0,1}$, $(A_{1,1}+2A)^2_{s,0,-1}$	$k < 0, \quad 0 < s$
C_p^5 $p \geq 0$	$I_{(1,2)}$	
	$(A_{1,1}+2A)^1_{k,0,1}$, $(A_{1,1}+2A)^1_{s,0,-1}$	$0 < k, \quad s < 0$
	$(A_{1,1}+2A)^2_{k,0,1}$, $(A_{1,1}+2A)^2_{s,0,-1}$	$k < 0, \quad 0 < s$
$(A_{1,1}+2A)^1$	$I_{(1,2)}$	
$(A_{1,1}+2A)^2$	$I_{(1,2)}$	

For three dimensional dual Lie superalgebras $(A_{1,1}+2A)^1_{\alpha,\beta,\gamma}$ and $(A_{1,1}+2A)^2_{\alpha,\beta,\gamma}$ which are isomorphic with $(A_{1,1}+2A)^1$ and $(A_{1,1}+2A)^2$, respectively, we have the following commutation relations:

$$\{\tilde{X}^2, \tilde{X}^2\} = i\alpha\tilde{X}^1, \quad \{\tilde{X}^2, \tilde{X}^3\} = i\beta\tilde{X}^1, \quad \{\tilde{X}^3, \tilde{X}^3\} = i\gamma\tilde{X}^1, \qquad \alpha, \beta, \gamma \in \Re. \tag{27}$$

Note that these Lie superalgebras are non isomorphic and they differ in the bound of their parameters.

Meanwhile note that for every Lie superalgebra of tables 1 and 2 there are trivial solutions for Eqs. (14) and (15) with $\tilde{f}^{ij}{}_k = 0$. (i.e. the dual Lie superalgebra is Abelian) we denote these dual Lie superalgebras with

$I_{(m,n)}$.

We see that for two and three dimensional Lie super-bialgebras (with two fermions), trivial Lie superalgebras are only dual to nontrivial one. The solution of super Jacobi and mixed super Jacobi identities and related isomorphism matrices C (in step 1) are listed in Appendix B. Here for explanation of steps 1-3 of section 3 we give an example.

5.1 An example

What follows is an explanation of the details of calculations for obtaining dual Lie superalgebras of Lie super-bialgebras $(C^4, (A_{1,1}+2A)^1_{k,0,1})$ and $(C^4, (A_{1,1}+2A)^1_{s,0,-1})$. As mentioned in Appendix B, the solution of super Jacobi and mixed super Jacobi identities has the following form:

$$\tilde{f}^{22}{}_1 = i\alpha, \qquad \tilde{f}^{33}{}_1 = i\beta, \qquad \tilde{f}^{23}{}_1 = i\gamma, \quad \forall \alpha, \beta, \gamma \in \Re. \tag{28}$$

This solution is isomorphic with the Lie superalgebra $(A_{1,1}+2A)^1$ with the following matrix:

$$C_1 = \begin{pmatrix} c_{11} & 0 & 0 \\ c_{21} & c_{22} & c_{23} \\ c_{31} & 0 & c_{33} \end{pmatrix}, \tag{29}$$

where $c_{11} = -ic_{33}^2\tilde{f}^{33}{}_1$, $c_{23} = -c_{22}\frac{\tilde{f}^{23}{}_1}{\tilde{f}^{33}{}_1}$, $\tilde{f}^{22}{}_1 = (\frac{c_{23}^2+c_{33}^2}{c_{22}^2})\tilde{f}^{33}{}_1$ and c_{22}, $c_{33} \in \Re - \{0\}$. By imposing that C_1 must be the transformation matrix (38), we have $c_{31} = c_{21} = 0$.
Now with the help of automorphism supergroup of C^4, the solution of (21) for the matrix B will be

$$B = \begin{pmatrix} \frac{c^2(b_{32}^2+b_{33}^2)}{\beta} & 0 & 0 \\ 0 & -\frac{b_{33}c_{33}}{c_{22}} & \frac{b_{32}c_{33}}{c_{22}} \\ 0 & b_{32} & b_{33} \end{pmatrix}, \quad \beta = -i\tilde{f}^{33}{}_1. \tag{30}$$

Now using (20), we obtain the following commutation relations for the dual Lie superalgebra $(A_{1,1}+2A)^{1\prime}$:

$$\{\tilde{X}^2, \tilde{X}^2\} = ia^{'}\tilde{X}^1, \qquad \{\tilde{X}^3, \tilde{X}^3\} = ib^{'}\tilde{X}^1, \qquad a^{'} = \frac{\beta c_{33}^2}{c^2 c_{22}^2}, \;\; b^{'} = \frac{\beta}{c^2}, \tag{31}$$

such that $a^{'}$ and $b^{'}$ have the same sings.
One cannot choose $a^{'} = b^{'}$ because in this case we have

$$B_1 = \begin{pmatrix} \frac{c^2({b^{'}}^2_{32}+{b^{'}}^2_{33})}{\beta} & 0 & 0 \\ 0 & -b^{'}_{33} & b^{'}_{32} \\ 0 & b^{'}_{32} & b^{'}_{33} \end{pmatrix}, \tag{32}$$

so as $BB_1^{-1} \notin A^{st}(C^4)$. Now by choosing $b^{'} = 1$ i.e., $\beta = c^2$ we have

$$B_2 = \begin{pmatrix} {b^{''}}^2_{32} + {b^{''}}^2_{33} & 0 & 0 \\ 0 & -\frac{c^{''}_{33}}{c^{''}_{22}}b^{''}_{33} & \frac{c^{''}_{33}}{c^{''}_{22}}b^{''}_{32} \\ 0 & b^{''}_{32} & b^{''}_{33} \end{pmatrix}. \tag{33}$$

such that $BB_2^{-1} \in A^{st}(C^4)$, this means that we can choose $b^{'} = 1$ and in this case, we have the following dual Lie superalgebras ($a^{'} = \frac{c_{33}^2}{c_{22}^2} = k > 0$):

$$(A_{1,1}+2A)^1_{k,0,1}: \qquad \{\tilde{X}^2, \tilde{X}^2\} = ik\tilde{X}^1, \qquad \{\tilde{X}^3, \tilde{X}^3\} = i\tilde{X}^1, \;\; k > 0. \tag{34}$$

In the same way by choosing $b^{'}=-1$ i.e., $\beta=-c^2$ such that $a^{'}\neq b^{'}$ we have

$$B_3=\begin{pmatrix} -(b^{'''2}_{\;\;32}+b^{'''2}_{\;\;33}) & 0 & 0\\ 0 & -\frac{c^{'''}_{33}}{c^{'''}_{22}}b^{'''}_{33} & \frac{c^{'''}_{33}}{c^{'''}_{22}}b^{'''}_{32}\\ 0 & b^{'''}_{32} & b^{'''}_{33}\end{pmatrix}. \tag{35}$$

so as $BB_3^{-1}\in A^{st}(C^4)$, this means that we can choose $b^{'}=-1$ and in this case, we have the following dual Lie superalgebras ($a^{'}=-\frac{c_{33}^2}{c_{22}^2}=s<0$):

$$(A_{1,1}+2A)^1_{s,0,-1}: \qquad \{\tilde{X}^2,\tilde{X}^2\}=is\tilde{X}^1, \qquad \{\tilde{X}^3,\tilde{X}^3\}=-i\tilde{X}^1, \;\; s<0. \tag{36}$$

Note that as $B_2B_3^{-1}\notin A^{st}(C^4)$, this means that the Lie super-bialgebras $(C^4,(A_{1,1}+2A)^1_{k,0,1})$ and $(C^4,(A_{1,1}+2A)^1_{s,0,-1})$ are not equivalent.

6 Conclusion

We have presented a new method for obtaining and classifying low dimensional *Lie super-bialgebras* which can also be applied for classification of low dimensional *Lie bialgebras*. We have classified all of the 48 two and three dimensional Lie super-bialgebras [22]. Using this classification one can construct Poisson-Lie T-dual sigma models [23] on the low dimensional Lie supergroups [24] (as Ref. [4] for Lie groups). Moreover, one can determinate the coboundary type of these Lie super-bialgebras and their classical r-matrices and Poisson brackets for their Poisson-Lie supergroups [25]. Determination of doubles and their isomorphism and investigation of Poisson-Lie plurality [26] are open problems for further investigation.

Acknowledgments

We would like to thank S. Moghadassi and F. Darabi for carefully reading the manuscript and useful comments.

Appendix A: Some properties of matrices and tensors in supervector spaces

In this appendix, we will review some basic properties of tensors and matrices in Ref. [16].

1. We consider the standard basis for the supervector spaces so that in writing the basis as a column matrix, we first present the bosonic base, then the fermionic one. The transformation of standard basis and its dual bases can be written as follows:

$$e'_i=(-1)^j K_i{}^j e_j, \qquad e'^i=K^{-st\,i}{}_j\, e^j, \tag{37}$$

where the transformation matrix K has the following block diagonal representation [16]

$$K=\left(\begin{array}{c|c} \mathrm{A} & \mathrm{C}\\ \hline \mathrm{D} & \mathrm{B}\end{array}\right), \tag{38}$$

where A, B and C are real submatrices and D is pure imaginary submatrix [16]. Here we consider the matrix and tensors having a form with all upper and lower indices written in the right hand side.

2. The transformation properties of upper and lower right indices to the left one for general tensors are as follows:

$$^iT^k_{jl\ldots}=T^{ik}_{jl\ldots}, \qquad {}_jT^{ik}_{l\ldots}=(-1)^j\,T^{ik}_{jl\ldots}. \tag{39}$$

3. For supertransposition we have

$$L^{st\,i}{}_j=(-1)^{ij}\,L_j{}^i, \qquad L^{st\;j}_{i}=(-1)^{ij}\,L^i{}_j,$$

$$M^{st}_{ij} = (-1)^{ij}\, M_{ji}, \qquad M^{st\, ij} = (-1)^{ij}\, M^{\,ji}. \tag{40}$$

4. For superdeterminant we have

$$sdet\left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array}\right) = det(A - CB^{-1}D)(detB)^{-1}, \tag{41}$$

when $detB \neq 0$ and

$$sdet\left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array}\right) = (det(B - DA^{-1}C))^{-1}\, (detA), \tag{42}$$

when $detA \neq 0$. For the inverse of matrix we have

$$\left(\begin{array}{c|c} \mathrm{A} & \mathrm{C} \\ \hline \mathrm{D} & \mathrm{B} \end{array}\right)^{-1} = \left(\begin{array}{c|c} (1_m - A^{-1}CB^{-1}D)^{-1}A^{-1} & -(1_m - A^{-1}CB^{-1}D)^{-1}A^{-1}CB^{-1} \\ \hline -(1_n - B^{-1}DA^{-1}C)^{-1}B^{-1}DA^{-1} & (1_n - B^{-1}DA^{-1}C)^{-1}B^{-1} \end{array}\right), \tag{43}$$

where $detA, detB \neq 0$ and m,n are dimensions of submatrices A and B, respectively.

Appendix B: Solutions $(sJ - msJ)$ for dual Lie superalgebras and isomorphism matrices

This appendix includes solutions of super Jacobi and mixed super Jacobi identities $(sJ - msJ)$ for dual Lie superalgebras and isomorphism matrices C which relate these solutions to other Lie superalgebras.

1. Solution of $(sJ - msJ)$ for dual Lie superalgebras of B is

$$\tilde{f}^{22}{}_{1} = i\alpha, \qquad \tilde{f}^{12}{}_{1} = 0, \tag{44}$$

where $\alpha \in \Re$.
Isomorphism matrix between these solutions and $(A_{1,1} + A)$ is as follows:

$$C = \left(\begin{array}{cc} -ic_{22}^2\tilde{f}^{22}{}_{1} & 0 \\ c_{21} & c_{22} \end{array}\right), \tag{45}$$

with the conditions $c_{22} \in \Re - \{0\}$ and imposing that C must be the transformation matrix, we have $c_{21} = 0$.

2. Solutions of $(sJ - msJ)$ for dual Lie superalgebras of C^1_p $(p \in \Re - \{0\})$ are

$i)$ $\quad \tilde{f}^{12}{}_{1} = \alpha, \;\; \tilde{f}^{23}{}_{3} = p\alpha, \qquad p \in \Re - \{0\}, \quad \alpha \in \Re,$

$ii)$ $\quad \tilde{f}^{12}{}_{1} = \tilde{f}^{23}{}_{3} = \beta, \qquad p = 1, \quad \beta \in \Re,$

$iii)$ $\quad \tilde{f}^{33}{}_{1} = i\gamma, \qquad p \in \Re - \{0\}, \quad \gamma \in \Re,$

$iv)$ $\quad \tilde{f}^{33}{}_{1} = i\lambda, \;\; \tilde{f}^{33}{}_{2} = i\eta \qquad p = \frac{1}{2}, \quad \lambda, \eta \in \Re,$

$v)$ $\quad \tilde{f}^{12}{}_{1} = \mu, \;\; \tilde{f}^{23}{}_{3} = -\frac{\mu}{2}, \;\; \tilde{f}^{33}{}_{1} = i\nu \qquad p = -\frac{1}{2}, \quad \mu, \nu \in \Re.$

For solution (i) we have isomorphism matrix between these solutions and C^1_p as follows:

$$C = \left(\begin{array}{ccc} c_{11} & -\frac{1}{\tilde{f}^{12}{}_{1}} & c_{13} \\ c_{21} & 0 & 0 \\ 0 & 0 & c_{33} \end{array}\right), \tag{46}$$

with the conditions $\tilde{f}^{23}{}_{3} = p\tilde{f}^{12}{}_{1}$ and $c_{21}, c_{33} \in \Re - \{0\}$, $c_{11} \in \Re$, as well as imposing that C must be transformation matrix we have $c_{13} = 0$. Solution (ii) is a special case of solution (i). For solution (iii) and (iv) $sdetC = 0$. Solution (v) is investigated for $(C^1_{\frac{1}{2}}, C^1_p)$ in the following step.

3. Solutions of $(sJ - msJ)$ for dual Lie superalgebras of $C^1_{\frac{1}{2}}$ are

$i)$ $\quad \tilde{f}^{23}{}_{3} = \alpha, \;\; \tilde{f}^{12}{}_{1} = 2\alpha, \quad \alpha \in \Re,$

$ii)$ $\quad \tilde{f}^{33}{}_{1} = i\beta, \;\; \tilde{f}^{33}{}_{2} = i\gamma, \;\; \tilde{f}^{23}{}_{3} = \frac{\beta}{2}, \;\; \tilde{f}^{13}{}_{3} = \frac{-\gamma}{2}, \;\; \tilde{f}^{12}{}_{1} = -\beta, \;\; \tilde{f}^{12}{}_{2} = -\gamma, \qquad \beta, \gamma \in \Re.$

For solution (i) we have isomorphism matrix C between $(C^1_{\frac{1}{2}}, C^1_p)$ as follows:

$$C = \left(\begin{array}{ccc} c_{11} & -\frac{1}{2\tilde{f}^{23}{}_3} & c_{13} \\ c_{21} & 0 & 0 \\ 0 & 0 & c_{33} \end{array} \right), \tag{47}$$

with the conditions $\tilde{f}^{12}{}_3 = 2\tilde{f}^{23}{}_3$, c_{21}, $c_{33} \in \Re - \{0\}$, $c_{11} \in \Re$ and $p = -\frac{1}{2}$, as well as imposing that C must be the transformation matrix, we have $c_{13} = 0$.

For solution (ii) we have isomorphism matrices C_1 and C_2 between $(C^1_{\frac{1}{2}}, C^1_{\frac{1}{2}})$ as follows:

$$C_1 = \left(\begin{array}{ccc} c_{11} & \frac{i}{\tilde{f}^{33}{}_1} & 0 \\ -ic_{33}^2\tilde{f}^{33}{}_1 & 0 & 0 \\ 0 & 0 & c_{33} \end{array} \right), \tag{48}$$

with the conditions $\tilde{f}^{33}{}_2 = \tilde{f}^{13}{}_3 = \tilde{f}^{12}{}_2 = 0$, $c_{33} \in \Re - \{0\}$ and $c_{11} \in \Re$

$$C_2 = \left(\begin{array}{ccc} c_{12}\frac{\tilde{f}^{33}{}_1}{\tilde{f}^{33}{}_2} - \frac{i}{\tilde{f}^{33}{}_2} & c_{12} & 0 \\ -ic_{33}^2\tilde{f}^{33}{}_1 & -ic_{33}^2\tilde{f}^{33}{}_2 & 0 \\ 0 & 0 & c_{33} \end{array} \right), \tag{49}$$

with the conditions $c_{33} \in \Re - \{0\}$ and $c_{12} \in \Re$.

4. Solution of $(sJ - msJ)$ for dual Lie superalgebras of C^2_p, C^3, C^4 and C^5_p is

$$\tilde{f}^{22}{}_1 = i\alpha, \qquad \tilde{f}^{33}{}_1 = i\beta, \qquad \tilde{f}^{23}{}_1 = i\gamma, \tag{50}$$

where α,β and $\gamma \in \Re$. Furthermore for C^3 we have another solution as follows:

$$\tilde{f}^{22}{}_1 = i\alpha, \qquad \alpha \in \Re,$$

where for this solution we have $sdetC = 0$, therefore we omit it.

4.1. For the above solution, we have the following isomorphism matrices C_1, C_2 and C_3 between C^2_p, C^3, C^4 and C^5_p with $(A_{1,1} + 2A)^1$:

$$C_1 = \left(\begin{array}{ccc} -ic_{33}^2\tilde{f}^{33}{}_1 & 0 & 0 \\ c_{21} & c_{22} & -c_{22}\frac{\tilde{f}^{23}{}_1}{\tilde{f}^{33}{}_1} \\ c_{31} & 0 & c_{33} \end{array} \right), \tag{51}$$

with the conditions $\tilde{f}^{22}{}_1 = (\frac{c_{23}^2 + c_{33}^2}{c_{22}^2})\tilde{f}^{33}{}_1$ and c_{22}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_1 must be the transformation matrix, we have $c_{31} = c_{21} = 0$,

$$C_2 = \left(\begin{array}{ccc} -ic_{23}^2\tilde{f}^{33}{}_1 & 0 & 0 \\ c_{21} & 0 & c_{23} \\ c_{31} & -c_{33}\frac{\tilde{f}^{33}{}_1}{\tilde{f}^{23}{}_1} & c_{33} \end{array} \right), \tag{52}$$

with the conditions $\tilde{f}^{22}{}_1 = (\frac{c_{23}^2 + c_{33}^2}{c_{32}^2})\tilde{f}^{33}{}_1$ and c_{23}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_2 must be transformation matrix, we have $c_{31} = c_{21} = 0$,

$$C_3 = \left(\begin{array}{ccc} c_{11} & 0 & 0 \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{array} \right), \tag{53}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{23}^2+c_{33}^2}{c_{22}^2+c_{32}^2})\tilde{f}^{33}{}_{1}$, $\tilde{f}^{23}{}_{1} = -(\frac{c_{33}c_{32}+c_{22}c_{23}}{c_{22}^2+c_{32}^2})\tilde{f}^{33}{}_{1}$ and $c_{11} \in \Re - \{0\}$, c_{22}, c_{23}, c_{32}, $c_{33} \in \Re$ with the conditions $c_{23}^2 + c_{33}^2 \neq 0$, $c_{22}^2 + c_{32}^2 \neq 0$, as well as imposing that C_3 must be the transformation matrix, we have $c_{31} = c_{21} = 0$.

4.2. For the above solution we have isomorphism matrices C_4, C_5, C_6, C_7 and C_8 between C_p^2, C^3, C^4 and C_p^5 with $(A_{1,1} + 2A)^2$ as follows:

$$C_4 = \begin{pmatrix} -ic_{32}(c_{23}-c_{33})\tilde{f}^{23}{}_{1} & 0 & 0 \\ c_{21} & c_{32} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{pmatrix}, \tag{54}$$

with the conditions $\tilde{f}^{22}{}_{1} = -(\frac{c_{23}+c_{33}}{c_{32}})\tilde{f}^{23}{}_{1}$, $\tilde{f}^{33}{}_{1} = 0$, $c_{23} \neq c_{33}$, $c_{32} \in \Re - \{0\}$ and $c_{23}, c_{33} \in \Re$, as well as imposing that C_4 must be the transformation matrix, we have $c_{31} = c_{21} = 0$,

$$C_5 = \begin{pmatrix} ic_{32}(c_{23}+c_{33})\tilde{f}^{23}{}_{1} & 0 & 0 \\ c_{21} & -c_{32} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{pmatrix}, \tag{55}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{23}-c_{33}}{c_{32}})\tilde{f}^{23}{}_{1}$, $\tilde{f}^{33}{}_{1} = 0$, $c_{23} \neq -c_{33}$, $c_{32} \in \Re - \{0\}$ and $c_{23}, c_{33} \in \Re$, as well as imposing that C_5 must be the transformation matrix, we have $c_{31} = c_{21} = 0$,

$$C_6 = \begin{pmatrix} -ic_{23}^2\tilde{f}^{33}{}_{1} & 0 & 0 \\ c_{21} & 0 & c_{23} \\ c_{31} & -c_{33}\frac{\tilde{f}^{33}{}_{1}}{\tilde{f}^{23}{}_{1}} & c_{33} \end{pmatrix}, \tag{56}$$

with the conditions $\tilde{f}^{22}{}_{1} = -(\frac{c_{23}^2-c_{33}^2}{c_{32}^2})\tilde{f}^{33}{}_{1}$ and c_{23}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_6 must be the transformation matrix, we have $c_{31} = c_{21} = 0$,

$$C_7 = \begin{pmatrix} ic_{33}^2\tilde{f}^{33}{}_{1} & 0 & 0 \\ c_{21} & c_{22} & -c_{22}\frac{\tilde{f}^{23}{}_{1}}{\tilde{f}^{33}{}_{1}} \\ c_{31} & 0 & c_{33} \end{pmatrix}, \tag{57}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{23}^2-c_{33}^2}{c_{22}^2})\tilde{f}^{33}{}_{1}$ and c_{22}, $c_{33} \in \Re - \{0\}$, as well as imposing that C_7 must be the transformation matrix, we have $c_{31} = c_{21} = 0$,

$$C_8 = \begin{pmatrix} c_{11} & 0 & 0 \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{pmatrix}, \tag{58}$$

with the conditions $\tilde{f}^{22}{}_{1} = (\frac{c_{33}^2-c_{23}^2}{c_{32}^2-c_{22}^2})\tilde{f}^{33}{}_{1}$, $\tilde{f}^{23}{}_{1} = (\frac{c_{22}c_{23}-c_{32}c_{33}}{c_{32}^2-c_{22}^2})\tilde{f}^{33}{}_{1}$ and $c_{11} \in \Re - \{0\}$, c_{22}, c_{23}, c_{32}, $c_{33} \in \Re$ with the conditions $c_{33}^2 - c_{23}^2 \neq 0, c_{32}^2 - c_{22}^2 \neq 0$, as well as imposing that C_8 must be the transformation matrix, we have $c_{31} = c_{21} = 0$.

References

[1] V. G. Drinfel'd, *Hamiltonian Lie groups, Lie bialgebras and the geometric meaning of the classical Yang-Baxter equation*, Sov. Math. Dokl. **27** (1983) 68-71.

[2] P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras,* I, Selecta Math. **2** (1996) no. 1, 1-41.
P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras*, II. Selecta Math. **4** (1998) no. 2, 213-231.

[3] P. Delorme, *Sur les triples de Manin pour les complexes,* Jour. Alg. **246** (2001) 97-174.

[4] M.A. Jafarizadeh and A. Rezaei-Aghdam, *Poisson-Lie T-duality and Bianchi type algebras*, *Phys. Lett. B* **458** (1999) 477-490, `[arXiv:hep-th/9903152]`.

[5] X. Gomez, *Classification of three-dimensional Lie bialgebras*, J. Math. Phys. **41** (2000) 4939.

[6] L. Hlavaty and L. Snobl, *Classification of 6-dimensional Manin triples*, `[arXiv:math-ph/0202209]`.

[7] A. Rezaei-Aghdam, M. Hemmati, A. R. Rastkar, *classification of real three-dimensional Lie bialgebras and their Poisson-Lie groups*, *J. Phys. A:Math.Gen.* **38** (2005) 3981-3994, `[arXiv:math-ph/0412092]`.

[8] Y. Kosmann-Schwarzbach, *Lie bialgebras, Poisson-Lie groups and dressing transformations integrability of nonlinear Systems*: Proc.(Pondicherry) ed Y Kosmann-Schwarzbach, B Grammaticos and K M Tamizhmani (Berlin: Springer, 1996) pp 104-70.

[9] N. Andruskiewitsch, *Lie Superbialgebras and Poisson-Lie supergroups*, Abh. Math. Sem. Univ. Hamburg **63** (1993) 147-163.

[10] N. Beisert and E. Spill, *The classical r-matrices of AdS/CFT and its Lie bialgebras structure*, `[arXiv:0708.1762 [hep-th]]`.

[11] N. Geer, *Etingof-Kazdan quantization of Lie super bialgebras*, `[arXiv:math.QA/0409563]`.

[12] C. Juszczak and J. T. Sobczyk, *Classification of low dimentional Lie super-bialgebras, J. Math. Phys.* **39**, (1998) 4982-4992.

[13] C. Juszczak, *Classification of* $osp(2|2)$ *Lie super-bialgebras,* `[arXiv:math.QA/9906101]`.

[14] G. Karaali, *A New Lie Bialgebra Structure on sl(2,1), Contemp. Math.* **413** (2006), pp.101-122.

[15] N. Backhouse, *A classification of four-dimensional Lie superalgebras*, *J. Math. Phys.* **19** (1978) 2400-2402.

[16] B. DeWitt, *Supermanifolds,* Cambridge University Press 1992.

[17] *Note that the bracket of one boson with one boson or one fermion is usual commutator but for one fermion with one fermion is anticommutator. Furthermore we identify grading of indices by the same indices in the power of (-1), for example grading*$(i) \equiv i$*; this is the notation that DeWitt applied in his book (Ref. [16]).*

[18] *For the proof of this proposition it is enough to use (11) and* $\delta'(X_i) = (-1)^{jk} \tilde{f}'^{jk}{}_{i} X_j \otimes X_k$ *and (22) in (5), then if super Jacobi and mixed super Jacobi identities [(2) and (12)] for* $\tilde{f}^{jk}{}_{i}$ *are satisfied then they satisfy for* $\tilde{f}'^{jk}{}_{i}$ *as well and vise versa.*

[19] A. Eghbali and A. Rezaei-Aghdam, *Classification of four-dimensional Lie super-bialgebras of the type* $(2,2)$, work in progress.

[20] *Note that the effect of A is linear and even i.e.*

$$A({a_i}^j X_j) = {a_i}^j A(X_j).$$

[21] *Note that the superalgebra A is one dimensional Abelian Lie superalgebras with one fermionic generator where Lie superalgebra* $A_{1,1}$ *is its bosonization.*

[22] *Note that in this classification one can omit the* $i = \sqrt{-1}$ *from commutation relations and to obtain Lie super-bialgebras in the nonstandard basis.*

[23] C. Klimčik and P. Ševera, *Dual non-Abelian duality and the Drinfeld double*, *Phys. Lett.* **B351** (1995) 445-462, `[arXiv:hep-th/9502122]`.
C. Klimčik, *Poisson-Lie T-duality, Nucl. Phys. Proc. Suppl.* **46** (1996) 116-121, `[arXiv:hep-th/9509095]`.

[24] A. Eghbali, A. Rezaei-Aghdam, *Poisson-Lie T-dual sigma models on supermanifolds*, JHEP 2009, **094** (2009), `[arXiv:0901.1592v3 [hep-th]]`.

[25] A. Eghbali and A. Rezaei-Aghdam, *Classical r-matrices of two and three dimensional Lie super-bialgebras and their Poisson-Lie supergroups*, [`arXiv:0908.2182 [math-ph]`].

[26] R. Von Unge, *Poisson-Lie T-plurality*, JHEP, 2002, **014** (2002) (preprint [`arXiv:hep-th/0205245`]).